

SCIENTIFIC REPORTS



OPEN

Long-term benefit of Microbiota Transfer Therapy on autism symptoms and gut microbiota

Dae-Wook Kang^{1,2,8}, James B. Adams³, Devon M. Coleman³, Elena L. Pollard³, Juan Maldonado^{1,2}, Sharon McDonough-Means⁴, J. Gregory Caporaso^{5,6} & Rosa Krajmalnik-Brown^{1,2,7}

Many studies have reported abnormal gut microbiota in individuals with Autism Spectrum Disorders (ASD), suggesting a link between gut microbiome and autism-like behaviors. Modifying the gut microbiome is a potential route to improve gastrointestinal (GI) and behavioral symptoms in children with ASD, and fecal microbiota transplant could transform the dysbiotic gut microbiome toward a healthy one by delivering a large number of commensal microbes from a healthy donor. We previously performed an open-label trial of Microbiota Transfer Therapy (MTT) that combined antibiotics, a bowel cleanse, a stomach-acid suppressant, and fecal microbiota transplant, and observed significant improvements in GI symptoms, autism-related symptoms, and gut microbiota. Here, we report on a follow-up with the same 18 participants two years after treatment was completed. Notably, most improvements in GI symptoms were maintained, and autism-related symptoms improved even more after the end of treatment. Important changes in gut microbiota at the end of treatment remained at follow-up, including significant increases in bacterial diversity and relative abundances of *Bifidobacteria* and *Prevotella*. Our observations demonstrate the long-term safety and efficacy of MTT as a potential therapy to treat children with ASD who have GI problems, and warrant a double-blind, placebo-controlled trial in the future.

The human gut and brain interact in complex ways, and abnormal conditions in the gut may predispose individuals to neurodevelopmental disorders^{1,2}. Individuals with Autism Spectrum Disorders (ASD)³, Parkinson's disease⁴, and Alzheimer's disease⁵, for example, have been known to experience chronic gastrointestinal (GI) symptoms as a common co-occurring medical condition, suggesting the presence of a gut-brain axis. Hallmayer *et al.*⁶ investigated 192 twin pairs and found that both genetic and environmental factors contribute to the etiology of ASD. The gut microbiome represents an important environmental factor that may exert an influence on symptoms, and a growing number of research groups have observed that children with ASD have distinctive gut microbiomes compared to neurotypical children^{7–11}. Moreover, multiple mouse studies have reported that gut microbes and their metabolites can impact behavior through the gut-brain axis, including for ASD^{12–14}.

Effective treatments for ASD include behavioral therapy, speech and social therapy, and dietary/nutritional/medical treatments, but no medical treatment has been approved to treat core symptoms of ASD¹⁵, such as social communication difficulties and repetitive behaviors. Considering the link between the gut and brain, modulating the gut microbiome by antibiotics, probiotics, prebiotics, and/or fecal microbiota transplant (FMT) could be a viable therapeutic option. In FMT, a large diversity and number of commensal microbes from a healthy donor are used to transform a dysbiotic gut microbiome into a healthy microbiome. In fact, FMT is the most effective therapy to treat recurrent *Clostridium difficile* infection¹⁶ and has shown varying levels of success for treating other

¹Biodesign Swette Center for Environmental Biotechnology, Arizona State University, Tempe, AZ, 85287, USA.

²Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University, Tempe, AZ, 85287, USA. ³School for Engineering of Matter, Transport and Energy, Arizona State University, Tempe, AZ, 85287, USA.

⁴Integrative Developmental Pediatrics, Tucson, AZ, 85701, USA. ⁵Center for Applied Microbiome Science, Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, AZ, 86001, USA. ⁶Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ, 86001, USA. ⁷School of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, AZ, 85287, USA. ⁸Present address: Department of Civil and Environmental Engineering, The University of Toledo, Toledo, OH, 43606, USA. Correspondence and requests for materials should be addressed to R.K.-B. (email: dr.rosy@asu.edu)




Figure 1. Changes in GI- and ASD-related symptoms of 18 children with ASD at two-year follow-up after treatment stopped. Asterisks (at the top of the box plot) indicate whether individuals (at each time point) have significantly changed since pre-treatment (Week 0 of original Phase 1 trial). Based on two-tailed Wilcoxon signed-rank test, *ns* indicates not significant, single asterisk indicates $p < 0.05$, double asterisks indicate $p < 0.01$, triple asterisks indicate $p < 0.001$. See also Supplementary Figs S1–S3.

GI disorders¹⁷, which has drawn attention to the method for use beyond GI-associated disorders¹⁸. Previously, we performed a pioneering open-label modified-FMT trial with an intensive combination called Microbial Transfer Therapy (MTT) consisting of two-week vancomycin treatment followed by a bowel cleanse and then high dose FMT for 1–2 days and 7–8 weeks of daily maintenance doses along with a stomach-acid suppressant, administered to children with ASD and chronic gastrointestinal problems¹⁹. After this 10-week MTT treatment and an eight-week follow-up observation period (18 weeks in total), we observed an 80% reduction in GI symptoms and a slow but steady improvement in core ASD symptoms. At the same time, we learned that gut microbial diversity, including potentially beneficial microbes, significantly increased after MTT¹⁹. Two years after this original clinical trial was completed, we re-evaluated the participants to determine whether observed improvements in behavior and GI symptoms persisted, and to ascertain the long-term impact of MTT on the gut microbiome of the study participants.

Results and Discussion

Improvements in GI and ASD symptoms remained two years after the MTT stopped.

Two years after the MTT was completed, we invited the 18 original subjects in our treatment group to participate in a follow-up study, and all provided informed consent. We performed the same GI and behavior tests that we employed previously¹⁹. 12 of 18 participants made some changes to their medication, diet, or nutritional supplements, but these changes were well documented and were mostly minor (Supplementary Table S1). We note that due to the open-label nature of this initial trial, all of the assessments are subject to placebo effect, however the long-term improvements we observed here are promising. Two years after treatment, most participants reported GI symptoms remaining improved compared to baseline (Fig. 1a and Supplementary Fig. S1). The improvement was on average 58% reduction in Gastrointestinal Symptom Rating Scale (GSRS) and 26% reduction in % days of abnormal stools (Daily Stool Record or DSR) relative to baseline, and this result is similar to what we observed at the end of treatment. The improvement in GI symptoms was observed for all sub-categories of GSRS (abdominal pain, indigestion, diarrhea, and constipation, Supplementary Fig. S2a) as well as for all sub-categories of DSR (no stool, hard stool, and soft/liquid stool, Supplementary Fig. S2b), although the degree of improvement on indigestion symptom (a sub-category of GSRS) was reduced after 2 years compared with weeks 10 and 18. This achievement is notable, because all 18 participants reported that they had had chronic GI problems (chronic constipation and/or diarrhea) since infancy, without any period of normal GI health (Supplementary Table S2). The families generally reported that ASD-related symptoms had slowly, steadily improved since week 18 of the Phase 1 trial, and this was consistent with the data reported in Fig. 1b–f. Based on the Childhood Autism Rating Scale (CARS) rated by a professional evaluator, the severity of ASD at the two-year follow-up was 47% lower than baseline (Fig. 1b), compared to 23% lower at the end of week 10. At the beginning of the open-label trial, 83% of participants rated in the severe ASD diagnosis per the CARS (Fig. 2a). At the two-year follow-up, only 17% were rated as severe, 39% were in the mild to moderate range, and 44% of participants were below the ASD diagnostic cut-off scores (Fig. 2a). The parent-rated Social Responsiveness Scale (SRS) assessment revealed that 89% of participants




Figure 2. CARS and SRS diagnostic category for ASD at baseline, 8 weeks after treatment, and two-year follow-up after treatment stopped. (a) For CARS, Minimal-to-No Symptoms (15–29.5 for ages less than 13; 15–27.5 for ages 13 or older), Mild-to-Moderate Symptoms (30–36.5 for ages less than 13; 28–34.5 for ages 13 or older), and Severe Symptoms (37 and higher for ages less than 13; 35 and higher for ages 13 or order)⁵⁴. (b) For SRS, Normal (0–53), Mild or High Functioning autism (54–86), Clinical diagnosis of autistic disorder, Asperger’s disorder, or more severe cases of Pervasive developmental disorder not otherwise specified (PDD-NOS) (>87)⁵⁵. (c–g) Strong and significant correlations between improvements in GI symptoms (GSRS) and behavior symptoms based on % changes in 2 years.

were in the severe range at the beginning of the trial, but the percentile dropped to 47% at the two-year follow-up (Fig. 2b), with 35% in the mild/moderate range and 18% below the cut-off for ASD. For the parent-rated Aberrant Behavior Checklist (ABC), total scores continued to improve, and were 35% lower relative to baseline (versus 24% lower at the end of treatment, relative to baseline; Fig. 1d). The Parent Global Impressions-III (PGI-III) scores remained similar to the scores at the end of treatment (week 10) of the open-label (Fig. 1e). The Vineland Adaptive Behavior Scale (VABS) equivalent age continued to improve (Fig. 1f), although not as quickly as during the treatment, resulting in an increase of 2.5 years over 2 years, which is much faster than typical for the ASD population, whose developmental age was only 49% of their physical age at the start of this study. Moreover, we observed improvement in behaviors in most sub-categories (Supplementary Figs S2c,d, and S3 for ABC, SRS, and VABS, respectively).

Overall, the most substantial improvements observed were on the CARS assessments, which was conducted by a professional evaluator and is less susceptible to placebo-effect²⁰. CARS is a stable and consistent diagnostic tool with high predictive validity²¹ and has been used to evaluate participants before and after therapeutic interventions in multiple studies^{20,22,23}. For the follow up CARS, the evaluator collected current information based on each question’s unique criteria. After the interview was complete for each question, the evaluator reviewed the information initially collected at baseline and used it for calibrating the final evaluation.




Figure 3. Stool microbiota assessments at two-year follow-up after treatment stopped. **(a)** Faith's phylogenetic diversity (PD) in the microbiota of 18 children with ASD as measured from stool samples. *Orange lines* indicate median PD of the donor samples (*dashed line* represents initial donor samples ($n = 5$), and *dotted line* represents maintenance dose samples ($n = 2$)), and *green line* indicates median PD of 20 neurotypical controls at week 0. *ns* indicates not significant, *single asterisk* indicates $p < 0.05$, *double asterisks* indicate $p < 0.01$, *triple asterisks* indicate $p < 0.001$ (two-tailed Wilcoxon signed-rank test comparing weeks 3, 10, and 18 and two-year to week 0 values). **(b)** Unweighted UniFrac distances between ASD gut microbiota and most relevant donor sample (initial donor sample at weeks 0 and 3, most recent maintenance dose sample at weeks 10 and 18, and 2 years). *Green line* indicates the median interpersonal variation between neurotypical controls and illustrates that prior to treatment the difference in gut microbiota composition between MTT recipients and donors was on the order of normal interpersonal variation. Statistics are the same as those used in (a). See also Supplementary Figs S5 and S6.

Improvements in GI and ASD symptoms were significantly correlated. We performed statistical analyses to assess whether improvements in GI and ASD severity were correlated. As shown in Fig. 2c–e, percentage changes in CARS, SRS, and ABC scores were positively correlated with percent changes in GRSR scores (Spearman correlation test, 2-tailed $p < 0.005$ and $r > 0.7$), implying that GI relief provided by MTT may ameliorate behavioral severity in children with ASD, or vice versa, or that both may be similarly impacted by another factor. Another GI assessment, DSR, however, showed that there was no significant correlation. Although the direction of the influence is not clear, a potential clinical link between GI and behavior severity is consistent with what previous studies have reported^{24,25}.

ASD fecal bacterial diversity was higher two years after the MTT stopped. 16 out of 18 original ASD participants provided an additional fecal sample two years after the open-label trial. Based on 16S ribosomal RNA (rRNA) gene amplicon sequencing analysis, most participants maintained higher gut microbiota diversity two years after treatment relative to baseline. Interestingly, for many individuals, the bacterial diversity was higher at two years than at the week 18 follow up as measured by Faith's Phylogenetic Diversity (Fig. 3a and Supplementary Fig. S4a) and Observed OTUs (Supplementary Fig. S5a). Considering low gut bacterial diversity in individuals with ASD²⁶ and other human disorders^{27–29}, an increase in diversity after MTT may reflect that MTT intervention successfully transformed gut environment into a healthier status and led to a long-term benefit on GI and behavior symptoms.

Upon completion of the original MTT treatment, we observed that the unweighted UniFrac distance³⁰ between the gut microbiota of MTT recipients and their corresponding donors was smaller than before treatment, suggesting some engraftment of the donor microbiome into the recipients by MTT¹⁹. Interestingly, two years after the trial, the recipients were as different from the donor microbiome as they were pre-treatment as measured by unweighted UniFrac distance (Fig. 3b, Supplementary Fig. S4b) and several other metrics of community dissimilarity (Supplementary Fig. S5b–e). This suggests that the recipients didn't retain completely the donated microbiome, but rather retained some features of it such as increased overall diversity, and increase in some important microbes such as *Prevotella*, while finding a new state.

Bifidobacterium and Prevotella relative abundances remained higher in feces of participants with ASD two-years after MTT stopped. Three taxa that were noticeably enhanced in MTT recipients at the conclusion of the original clinical trial¹⁹ were revisited during the two-year follow-up. Notably, compared to baseline, median relative abundances of *Bifidobacteria* and *Prevotella* increased 4-fold and 712-fold at week 10, and 5-fold and 84-fold at two years, respectively (Fig. 4a,b). *Desulfovibrio* relative abundance decreased since week 18 (Fig. 4c), but at the two-year follow-up was still marginally higher compared to baseline (two-tailed Wilcoxon signed-rank test, $p = 0.07$) and higher than neurotypical controls (two-tailed Mann-Whitney U test, $p < 0.05$). An increase in *Prevotella* after MTT is noteworthy, since its lower abundance in feces of children with ASD compared with neurotypical children has been confirmed in two different cohorts^{26,31}. A recent study also found reduced levels of *Prevotella* in the oral microbiome of children with ASD³². *Prevotella* may be involved in butyrate production³³, a key nutrient for the intestinal epithelial cells³⁴. In addition, its co-occurrence with *Desulfovibrio* may reflect a synergistic advantage to outcompete other commensal microbes that utilize mucin as nutrients³⁵, although more research is needed on how their ecological niche in mucin desulfation could




Figure 4. Changes in relative abundances of *Bifidobacterium*, *Prevotella*, and *Desulfovibrio*. *ns* indicates not significant, *single asterisk* indicates $p < 0.05$ and *double asterisks* indicate $p < 0.01$ (two-tailed Wilcoxon signed-rank test comparing weeks 3, 10, and 18 and two-year to week 0 values). *Orange lines* indicate median of the donor samples (*dashed line* represents initial donor samples, and *dotted line* represents maintenance dose samples), and *green line* indicates median of 20 neurotypical controls at week 0.

contribute to an integrity of gut epithelial cells³⁶ as well as to the improvement on GI and behavior symptoms we observed. Further mechanistic studies with multi-omic approaches are warranted to define the roles of *Prevotella* and *Desulfovibrio* in the context of autism.

Further research with a placebo-control arm is warranted. To the best of our knowledge, long-term follow-up studies are rare for medical treatment of individuals with ASD. In treatments with vancomycin³⁷ or phytochemical sulforaphane³⁸, benefits were lost within two or four weeks, respectively, of the treatments being discontinued. Thus, the long-term benefits observed here two years after MTT stopped are very encouraging, and MTT-driven gut microbiota transformation seems robust and long-lasting for the treatment of ASD. Despite steady and continuous improvement in behaviors over two years, we must underscore that the original clinical trial and current follow-up study are open-label trials without a control for placebo effect. Autism symptoms are relatively stable over time without a major intervention: for example, a trajectory study with 345 children with ASD showed that more than 80% of participants with ASD retained unexpectedly stable core symptoms severity over 8 to 12 years³⁹. The VABS observations indicate that the improvements in adaptive behaviors observed here were substantially more than expected for children with ASD over two years. A limitation of this study is that 12 out of 18 participants made one or more changes to their medications, nutritional supplements, and diets between the end of the original MTT trial and the two-year follow-up since the treatment stopped (Supplementary Table S1). As described in detail in the methods section, participants were asked to rate the perceived effectiveness on GI and ASD symptoms (on a scale of 0–4) caused by changes in medications, diet, or nutritional supplements. Although the scale on the perceived effectiveness is still subjective and difficult to interpret, low scores received (1.1 for GI and 0.8 for ASD symptoms) suggest that these treatments on average could have only “slight effect”. Thus, it appears that most of the changes observed were probably due to the MTT, although we still need follow-up studies to understand whether the improvement by MTT were solely from vancomycin, MoviPrep, Prilosec, Standardized Human Gut Microbiota (SHGM), or a combination of these individual factors. For example, some participants in our study could have GI symptoms that were acid-peptic in nature, and their improvements on GI symptoms might be solely attributed to the administration of stomach-acid suppressant (Prilosec)⁴⁰. We hypothesize that MTT may also be beneficial for children with ASD who do not have obvious GI symptoms but have low diversity of gut bacteria, as our previous study²⁶ found that most children with ASD had low gut bacterial diversity, regardless of whether they have GI problems.

Here, we also would like to address a potential study limitation interpreting the improvement on GI symptoms after MTT, since these heterogeneous GI symptoms could reflect a wide range of underlying etiological GI pathologies. Although we reviewed medical histories to exclude children with known gastrointestinal diagnoses (such as ulcerative colitis, Crohn’s Disease, celiac disease, eosinophilic gastritis, or similar conditions)¹⁹, we did not conduct additional GI diagnostic evaluations, which is a limitation of this study. Thus, we want to underscore need of follow-up studies embracing more thorough examination of participants’ GI pathologies in order to better understand effectiveness of MTT.

Conclusions

In summary, all 18 participants with ASD were re-evaluated two years after MTT treatment stopped, and we observed significant improvements both in GI and behavior symptoms as compared with baseline measurements collected at the beginning of the original open-label trial. GI benefits were mostly maintained from the end of treatment, and autism symptoms were reported to have improved significantly since the end of treatment. Changes in gut microbiota persisted at two years, including in overall community diversity and relative abundances of *Bifidobacteria* and *Prevotella*. These encouraging observations demonstrate that the intensive MTT

intervention is a promising therapy for treating children with ASD who have GI problems. We recommend future research including double-blind, placebo-controlled randomized trials with a larger cohort.

Methods

Ethics approval and consent to participate. The protocol for the original treatment study was approved by FDA (Investigational new drug number 15886) and the Institutional Review Board of Arizona State University (ASU IRB Protocol #: 00001053), as described in Kang *et al.*¹⁹. All research methods were performed in accordance with the relevant guidelines and regulations. For the follow-up study described in this paper, a new IRB protocol was approved by the IRB of Arizona State University (ASU IRB Protocol # 00004890). The follow-up study was only open to the 18 children with ASD who participated in the original Phase 1 trial study, and no further inclusion or exclusion criteria was applied to what we employed in the original trial¹⁹. For the consent process, we contacted past participants and their parents and explained the study in detail by phone and/or email, and provided them with a copy of the parent permission form and the child assent form. After the informed consent process, all 18 families agreed to participate and signed written parent permission and assent forms. We maintained confidentiality of participants results by de-identifying all samples for the entire analyses. The participant's name and identifiers were removed and not used in all sections of the manuscript, including supplementary information. The trial was registered at the ClinicalTrials.gov (NCT02504554) on March 30, 2015.

Study objective. The purpose of the study was to conduct a 2-year follow-up on children with ASD who participated in the original study and to determine long-term safety and efficacy of the Microbial Transfer Therapy (MTT). In brief, the previous study was designed to perform an open-label clinical trial for children with ASD and moderate to severe GI problems (ages 7–17 years)¹⁹. Eighteen children were enrolled in the previous study and treated with a combination of vancomycin (to reduce pathogenic bacteria) for 2 weeks, MoviPrep (a bowel cleanse to remove vancomycin and remaining bacteria), Prilosec (a proton pump inhibitor in order to reduce stomach acidity and to increase the survival rate of high and maintenance doses of SHGM), and Standardized Human Gut Microbiota (gut bacteria from healthy donors) for 7–8 weeks⁴¹. Participants were monitored during the 10 weeks of treatment, and again 8 weeks after MTT stopped. There were no major adverse effects, and all 18 participants completed the study. Most of the children had substantial reduction of their GI and ASD symptoms, and those improvements remained at the follow-up at 8 weeks after the treatment ended. Further information on the previous study design and outcomes are described in Kang *et al.*¹⁹. In this current study, we followed up past participants approximately 2 years since the treatment stopped, and requested parents to collect one stool sample from their child in order to investigate changes in bacterial composition. After the study ended, 6 participants did not make any changes in their treatments for the next two years, and 12 participants made one or more changes (Supplementary Table S1). The changes (and the number of participants making those changes) included dietary changes (gluten-free/casein-free diet (1), ketogenic diet (1), and removing eggs(1)), GI treatments (magnesium citrate (2), digestive enzymes (1), dulcolax (1), enteragum (1), probiotics (2)), medication changes (Abilify (1), birth control (1), IVIG (1), Lamictal (1), lithium (1), Tenex (1), thyroid (1), and Vyvanse (1), and nutritional supplements (amino acids (1), multivitamins (2), Bravo yogurt (1), citruline (1), fish oil (1), GABA (1), l-theanine (1), magnesium (1), Oraimmune (1), and Rerum (1)). Participants were asked to rate the perceived effectiveness of each treatment on a scale of 0–4 for both GI and ASD symptoms, where 0 = “No effect”, 1 = “Slight benefit”, 2 = “Moderate benefit”, 3 = “Good benefit”, and 4 = “Great benefit”.

Assessments of gastrointestinal and autism-related symptoms. To be consistent with the previous assessments¹⁹, we evaluated GI symptoms using a revised version of Gastrointestinal Symptom Rating Scale (GSRS) that employs 7-point Likert scale⁴² in 5 domains: Abdominal Pain, Reflux, Indigestion, Diarrhea, and Constipation. We also collected Daily Stool Records (DSR) that rates the stool using the Bristol Stool Form scale (1 = very hard, 7 = liquid) over 14 days. For the assessments of autism and related symptoms, we also employed the same assessments that we used in the previous study including the Parent Global Impressions–III (PGI-III)⁴³, the Childhood Autism Rating Scale (CARS), the Aberrant Behavior Checklist (ABC), the Social Responsiveness Scale (SRS), and the Vineland Adaptive Behavior Scale II (VABS-II). The CARS evaluation was conducted by the same professional evaluator who conducted the previous CARS evaluation, and parents assessed the GSRS, DSR, PGI-III, ABC, SRS, and VABS-II at approximately 2 years after the therapy stopped.

Microbial DNA extraction and next generation sequencing. In order to be consistent with the previous work in Kang *et al.*¹⁹, we employed the same DNA extraction assay, library preparation, and sequencing methods. We extracted microbial genomic DNA from stool samples using a PowerSoil[®] DNA Isolation Kit (Mobio Carlsbad, CA) and constructed 16S rRNA library for MiSeq Illumina platform according to the protocol from Earth Microbiome Project (<http://www.earthmicrobiome.org/protocols-and-standards/16s/>). The barcoded primer set 515f–806r was used for pair-ended sequencing to target the 16S rRNA V4 region⁴⁴. Library preparation and sequencing work were performed at the Microbiome Analysis Laboratory in the Bidesign Institute of Arizona State University (<http://krajmalnik.environmentalbiotechnology.org/microbiome-lab.html>).

Microbiome bioinformatics. The 16S rRNA gene data from our previous sequencing run was generated from four Illumina MiSeq runs, and the follow-up data (new to this study) was sequenced on a single MiSeq run. To ensure consistent processing across all of these runs, and to enable the use of the most recent bioinformatics methods for microbiome analysis, we processed all five of these sequencing runs together in a single QIIME 2 analysis⁴⁵.

Analyses were performed on the forward reads only, to retain as many sequences as possible (i.e., to not discard sequences where forward and reverse reads could not be joined). Sequence quality control was performed with two different processing pipelines, both implemented in QIIME 2.

The first processing pipeline used DADA2⁴⁶ via the q2-dada2 QIIME 2 plugin (we'll refer to this as the DADA2 pipeline). In the DADA2 pipeline, the first 10 bases were trimmed from reads and reads were truncated at 150 bases prior to processing with the dada2-single method, as recommended by the DADA2 authors. All other DADA2 parameters were left with their default settings. DADA2 was run on a per-sequencing run basis using the same parameters for all runs of DADA2. The resulting feature tables were merged, such that we had one master "feature table" containing counts of features on a per sample basis, where our features were unique 16S rRNA gene sequence variants. The "feature sequence" that defined each feature was the unique 16S rRNA gene sequence variant. Microbiome analysis results based on this processing pipeline are presented in Figs 3, 4 and Supplementary Figs S4, S5.

The second processing pipeline used the quality filtering approach described in Bokulich *et al.*⁴⁷ as implemented in the q2-quality-filter plugin using a min-quality of 20 (all other parameters were defaults). We refer to this pipeline as the Q20 pipeline. Sequence reads were subsequently clustered into 97% OTUs using open-reference OTU picking as implemented in the q2-vsearch plugin (which uses vsearch⁴⁸ for clustering). Again, we had one master "feature table" containing counts of features on a per sample basis, where in this case our features were 97% OTUs. The "feature sequence" that defined each feature was the OTU centroid sequence. As an additional quality control step, features that were only observed in a single sample and features that could not be assigned to a phylum against the Silva database⁴⁹ (described further below) were filtered from the feature table. Microbiome analysis results based on this processing pipeline are presented in Supplementary Fig. S6.

Following both the DADA2 and Q20 pipelines, feature sequences were aligned with mafft⁵⁰ (wrapped in the q2-alignment plugin), high entropy positions were filtered from the resulting alignment⁵¹, an unrooted tree was constructed with FastTree⁵² (wrapped in the q2-phylogeny plugin), and the tree was rooted using midpoint rooting. Taxonomy was assigned to each feature sequence against the Silva 119 database using a Naïve Bayes classifier implemented in the q2-feature-classifier plugin⁵³. Feature tables were rarefied to 5,298 sequences per sample, and diversity metrics were computed using the q2-diversity plugin.

For complete details on the bioinformatics methods, including the versions of all software and dependencies, and all commands and parameter settings, readers can view the provenance data contained in the Supplementary File S1. This information can be viewed visually at <https://view.qiime2.org>.

The specific values of the microbiome richness and distance metrics differ between the DADA2 and Q20 bioinformatics pipelines. For example, Faith's Phylogenetic Diversity is considerably lower with the DADA2 pipeline, likely because it does a much better job at raw sequence quality control (as described in the original DADA2 paper). Despite this, it is encouraging to observe that the different quality control and feature definition pipelines illustrate the same trends in microbiome composition and richness.

Statistical analysis. We assumed the data as non-normally distributed because of a relatively small sample size, and performed nonparametric analyses of Mann-Whitney U-test, Wilcoxon signed-rank test, and Spearman correlation test. We reported all *p* values from two-tailed tests, and accepted *p* values lower than 0.05 as significant. For the boxes in Figs 1, 3 and 4, the cross line in the box indicates median values and the bottom and top edges of the box indicate the 25th and 75th percentiles of the sample.

Data Availability

The 16S rRNA gene sequence reads are available in the open-source sequence data repository 'Qiita' with the study ID number 10532 (<https://qiita.ucsd.edu>) and the NCBI SRA under BioProject ID PRJNA529598. For details on obtaining and installing QIIME 2, see <https://qiime2.org>. QIIME 2 is open source and free for all use. The analyses presented in Fig. 3 and Supplementary Figs S4–S6 were performed using Jupyter Notebooks. These notebooks are available at <https://github.com/caporaso-lab/autism-fmt1>.

References

- Rogers, G. B. *et al.* From gut dysbiosis to altered brain function and mental illness: mechanisms and pathways. *Molecular Psychiatry* **21**, 738–748, <https://doi.org/10.1038/mp.2016.50> (2016).
- Sharon, G., Sampson, T. R., Geschwind, D. H. & Mazmanian, S. K. The central nervous system and the gut microbiome. *Cell* **167**, 915–932, <https://doi.org/10.1016/j.cell.2016.10.027> (2016).
- McElhanon, B. O., McCracken, C., Karpen, S. & Sharp, W. G. Gastrointestinal symptoms in autism spectrum disorder: a meta-analysis. *Pediatrics* **133**, <https://doi.org/10.1542/peds.2013-3995> (2014).
- Braak, H., Rub, U., Gai, W. P. & Del Tredici, K. Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *Journal of Neural Transmission* **110**, 517–536, <https://doi.org/10.1007/s00702-002-0808-2> (2003).
- McCarthy, M., AddingtonHall, J. & Altmann, D. The experience of dying with dementia: A retrospective study. *International Journal of Geriatric Psychiatry* **12**, 404–409, [https://doi.org/10.1002/\(SICI\)1099-1166\(199703\)12:3<404::AID-GPS529>3.0.CO;2-2](https://doi.org/10.1002/(SICI)1099-1166(199703)12:3<404::AID-GPS529>3.0.CO;2-2) (1997).
- Hallmayer, J. *et al.* Genetic heritability and shared environmental factors among twin pairs with autism. *Archives of General Psychiatry* **68**, 1095–1102, <https://doi.org/10.1001/archgenpsychiatry.2011.76> (2011).
- Finogold, S. M. *et al.* Pyrosequencing study of fecal microflora of autistic and control children. *Anaerobe* **16**, 444–453, <https://doi.org/10.1016/j.anaerobe.2010.06.008> (2010).
- De Angelis, M. *et al.* Fecal microbiota and metabolome of children with autism and pervasive developmental disorder not otherwise specified. *Plos One* **8**, 18, <https://doi.org/10.1371/journal.pone.0076993> (2013).
- Williams, B. L., Hornig, M., Parekh, T. & Lipkin, W. I. Application of novel PCR-based methods for detection, quantification, and phylogenetic characterization of *Sutterella* species in intestinal biopsy samples from children with autism and gastrointestinal disturbances. *MBio* **3**, e00261–11, <https://doi.org/10.1128/mBio.00261-11> (2012).
- Gondalia, S. V. *et al.* Molecular characterisation of gastrointestinal microbiota of children with autism (with and without gastrointestinal dysfunction) and their neurotypical siblings. *Autism Research* **5**, 419–427, <https://doi.org/10.1002/aur.1253> (2012).
- Son, J. S. *et al.* Comparison of fecal microbiota in children with autism spectrum disorders and neurotypical siblings in the Simons simplex collection. *Plos One* **10**, e0137725, <https://doi.org/10.1371/journal.pone.0137725> (2015).
- Hsiao, E. Y. *et al.* Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopment disorders. *Cell* **155**, 1451–1463, <https://doi.org/10.1016/j.cell.2013.11.024> (2013).

13. Kim, S. *et al.* Maternal gut bacteria promote neurodevelopmental abnormalities in mouse offspring. *Nature* **549**, 528–532, <https://doi.org/10.1038/nature23910> (2017).
14. Buffington, S. A. *et al.* Microbial reconstitution reverses maternal diet-induced social and synaptic deficits in offspring. *Cell* **165**, 1762–1775, <https://doi.org/10.1016/j.cell.2016.06.001> (2016).
15. Lacivita, E., Perrone, R., Margari, L. & Leopoldo, M. Targets for drug therapy for autism spectrum disorder: challenges and future directions. *Journal of Medicinal Chemistry* **60**, 9114–9141, <https://doi.org/10.1021/acs.jmedchem.7b00965> (2017).
16. Bagdasarian, N., Rao, K. & Malani, P. N. Diagnosis and treatment of *Clostridium difficile* in adults a systematic review. *Jama-Journal of the American Medical Association* **313**, 398–408, <https://doi.org/10.1001/jama.2014.17103> (2015).
17. Moayyedi, P. *et al.* Fecal microbiota transplantation induces remission in patients with active ulcerative colitis in a randomized controlled trial. *Gastroenterology* **149**, 102–109, <https://doi.org/10.1053/j.gastro.2015.04.001> (2015).
18. Xu, M. Q. *et al.* Fecal microbiota transplantation broadening its application beyond intestinal disorders. *World Journal of Gastroenterology* **21**, 102–111, <https://doi.org/10.3748/wjg.v21.i1.102> (2015).
19. Kang, D.-W. *et al.* Microbiota Transfer Therapy alters gut ecosystem and improves gastrointestinal and autism symptoms: an open-label study. *Microbiome* **5**, 10, <https://doi.org/10.1186/s40168-016-0225-7> (2017).
20. Belsito, K. M., Law, P. A., Kirk, K. S., Landa, R. J. & Zimmerman, A. W. Lamotrigine therapy for autistic disorder: A randomized, double-blind, placebo-controlled trial. *Journal of Autism and Developmental Disorders* **31**, 175–181, <https://doi.org/10.1023/a:1010799115457> (2001).
21. Nah, Y. H., Young, R. L. & Brewer, N. Using the Autism Detection in Early Childhood (ADEC) and Childhood Autism Rating Scales (CARS) to predict long term outcomes in children with autism spectrum disorders. *Journal of Autism and Developmental Disorders* **44**, 2301–2310, <https://doi.org/10.1007/s10803-014-2102-1> (2014).
22. Gattino, G. S., Riesgo, R. D., Longo, D., Leite, J. C. L. & Faccini, L. S. Effects of relational music therapy on communication of children with autism: a randomized controlled study. *Nordic Journal of Music Therapy* **20**, 142–154, <https://doi.org/10.1080/08098131.2011.566933> (2011).
23. Saad, K. *et al.* Randomized controlled trial of vitamin D supplementation in children with autism spectrum disorder. *Journal of Child Psychology and Psychiatry* **59**, 20–29, <https://doi.org/10.1111/jcpp.12652> (2018).
24. Adams, J. B., Johansen, L. J., Powell, L. D., Quig, D. & Rubin, R. A. Gastrointestinal flora and gastrointestinal status in children with autism—comparisons to typical children and correlation with autism severity. *Bmc Gastroenterology* **11**, 22, <https://doi.org/10.1186/1471-230x-11-22> (2011).
25. Horvath, K. & Perman, J. A. Autistic disorder and gastrointestinal disease. *Current Opinion in Pediatrics* **14**, 583–587, <https://doi.org/10.1097/01.mop.0000030221.71203.46> (2002).
26. Kang, D. W. *et al.* Reduced incidence of *Prevotella* and other fermenters in intestinal microflora of autistic children. *Plos One* **8**, e68322, <https://doi.org/10.1371/journal.pone.0068322> (2013).
27. Ilhan, Z. E. *et al.* Distinctive microbiomes and metabolites linked with weight loss after gastric bypass, but not gastric banding. *Isme Journal* **11**, 2047–2058, <https://doi.org/10.1038/ismej.2017.71> (2017).
28. Ahn, J. *et al.* Human gut microbiome and risk for colorectal cancer. *Jnci-Journal of the National Cancer Institute* **105**, 1907–1911, <https://doi.org/10.1093/jnci/djt300> (2013).
29. Ott, S. J. *et al.* Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active inflammatory bowel disease. *Gut* **53**, 685–693, <https://doi.org/10.1136/gut.2003.025403> (2004).
30. Lozupone, C. & Knight, R. UniFrac: a new phylogenetic method for comparing microbial communities. *Applied and Environmental Microbiology* **71**, 8228–8235, <https://doi.org/10.1128/aem.71.12.8228-8235.2005> (2005).
31. Kang, D. W. *et al.* Differences in fecal microbial metabolites and microbiota of children with autism spectrum disorders. *Anaerobe* **49**, 121–131, <https://doi.org/10.1016/j.anaerobe.2017.12.007> (2017).
32. Qiao, Y. A. *et al.* Alterations of oral microbiota distinguish children with autism spectrum disorders from healthy controls. *Scientific Reports* **8**, 12, <https://doi.org/10.1038/s41598-018-19982-y> (2018).
33. Esquivel-Elizondo, S., Ilhan, Z. E., Garcia-Pena, E. I. & Krajmalnik-Brown, R. Insights into butyrate production in a controlled fermentation system via gene predictions. *mSystems* **2**, 13, <https://doi.org/10.1128/mSystems.00051-17> (2017).
34. Donohoe, D. R. *et al.* The microbiome and butyrate regulate energy metabolism and autophagy in the mammalian colon. *Cell Metabolism* **13**, 517–526, <https://doi.org/10.1016/j.cmet.2011.02.018> (2011).
35. Arumugam, M. *et al.* Enterotypes of the human gut microbiome. *Nature* **473**, 174–180, <https://doi.org/10.1038/nature09944> (2011).
36. Derrien, M. *et al.* Mucin-bacterial interactions in the human oral cavity and digestive tract. *Gut Microbes* **1**, 254–268, <https://doi.org/10.4161/gmic.1.4.12778> (2010).
37. Sandler, R. H. *et al.* Short-term benefit from oral vancomycin treatment of regressive-onset autism. *Journal of Child Neurology* **15**, 429–435, <https://doi.org/10.1177/088307380001500701> (2000).
38. Singh, K. *et al.* Sulforaphane treatment of autism spectrum disorder (ASD). *Proceedings of the National Academy of Sciences of the United States of America* **111**, 15550–15555, <https://doi.org/10.1073/pnas.1416940111> (2014).
39. Gotham, K., Pickles, A. & Lord, C. Trajectories of autism severity in children using standardized ADOS scores. *Pediatrics* **130**, E1278–E1284, <https://doi.org/10.1542/peds.2011-3668> (2012).
40. Jain, K. S. *et al.* Recent advances in proton pump inhibitors and management of acid-peptic disorders. *Bioorganic & Medicinal Chemistry* **15**, 1181–1205, <https://doi.org/10.1016/j.bmc.2006.07.068> (2007).
41. Hamilton, M. J., Weingarden, A. R., Sadowsky, M. J. & Khoruts, A. Standardized frozen preparation for transplantation of fecal microbiota for recurrent *Clostridium difficile* infection. *Am J Gastroenterol* **107**, 761–767, <https://doi.org/10.1038/ajg.2011.482> (2012).
42. Revicki, D. A., Wood, M., Wiklund, I. & Crawley, J. Reliability and validity of the gastrointestinal symptom rating scale in patients with gastroesophageal reflux disease. *Quality of Life Research* **7**, 75–83, <https://doi.org/10.1023/A:1008841022998> (1997).
43. Adams, J. B. *et al.* Effect of a vitamin/mineral supplement on children and adults with autism. *BMC Pediatr* **11**, <https://doi.org/10.1186/1471-2431-11-111> (2011).
44. Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *Isme Journal* **6**, 1621–1624, <https://doi.org/10.1038/ismej.2012.8> (2012).
45. Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* **7**, 335–336, <https://doi.org/10.1038/nmeth.f.303> (2010).
46. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583, <https://doi.org/10.1038/nmeth.3869> (2016).
47. Bokulich, N. A. *et al.* Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nature Methods* **10**, 57–59, <https://doi.org/10.1038/nmeth.2276> (2013).
48. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahe, F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584, <https://doi.org/10.7717/peerj.2584> (2016).
49. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* **41**, D590–D596, <https://doi.org/10.1093/nar/gks1219> (2013).
50. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772–780, <https://doi.org/10.1093/molbev/mst010> (2013).
51. Lane, D. *Nucleic Acid Techniques in Bacterial Systematics*. 115–175 (John Wiley and Sons, 1991).

52. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2-approximately maximum-likelihood trees for large alignments. *Plos One* 5, e9490, <https://doi.org/10.1371/journal.pone.0009490> (2010).
53. Bokulich, N. A. *et al.* Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 6, 90, <https://doi.org/10.1186/s40168-018-0470-z> (2018).
54. Schopler, E., Van Bourgondien, M. E., Wellman, G. J. & Love, S. R. *Childhood autism rating scale, (CARS2)*. (WPS Torrance, CA, 2010).
55. Constantino, J. N. *Social Responsiveness Scale (SRS)*. (Western Psychological Services, 2005).

Acknowledgements

We gratefully thank all children with ASD and their families for participating in the follow-up study. We would like to thank Thomas Borody, Alexander Khoruts, Michael J. Sadowsky, and Alessio Fasano for their help with the treatment portion of the study and also for their help with the original Investigational New Drug (IND) application and study. We thank Finch Therapeutics (formerly Crestovo) for providing Standardized Human Gut Microbiota for the study. We also thank Carole Flores for editing assistance. This work was supported by Finch Therapeutics and the Arizona Board of Regents.

Author Contributions

D.-W.K., J.B.A. and R.K.B. designed the study, wrote funding proposal. D.-W.K., J.B.A., D.C. and coordinated the study. J.B.A., D.C., S.M.-M. and E.L.P. gathered the clinical data. D.-W.K., J.M. and J.G.C. processed the biosamples and performed the data acquisition. D.-W.K., J.B.A., J.M., D.C., J.G.C. and R.K.B. performed the analysis and interpreted the data. D.-W.K., J.B.A. and R.K.B. wrote the main paper, and J.G.C., J.M., S.M.-M., E.L.P., revised the manuscript. All authors read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-019-42183-0>.

Competing Interests: J.B.A., D.-W.K. and R.K.B. have pending/approved patents related to the use of FMT and/or probiotics for various conditions including autism (Application number: 14/403,425 (approved and active); 15/290,798 (pending)). J.B.A., R.K.B. and D.-W.K. have received research funding from Crestovo/Finch Therapeutics for FMT research. J.B.A., R.K.B. and J.G.C. have received consulting fees from Crestovo. The other authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019