



Article

Co-Administration of Iron and a Bioavailable Curcumin Supplement Increases Serum BDNF Levels in Healthy Adults

Helena Tiekou Lorinczova ^{1,†}, Owen Fitzsimons ^{1,†}, Leah Mursaleen ^{1,2}, Derek Renshaw ³, Gulshanara Begum ¹ and Mohammed Gulrez Zariwala ^{1,*}

¹ School of Life Sciences, University of Westminster, 115 New Cavendish Street, London W1W 6UW, UK; H.Lorinczova@westminster.ac.uk (H.T.L.); O.Fitzsimons@westminster.ac.uk (O.F.); w1655446@my.westminster.ac.uk (L.M.); begumru@westminster.ac.uk (G.B.)

² The Cure Parkinson's Trust, 120 New Cavendish St, Fitzrovia, London W1W 6XX, UK

³ Centre for Sport, Exercise and Life Sciences, Faculty of Health and Life Sciences, Coventry University, Priory St, Coventry CV1 5FB, UK; derek.renshaw@coventry.ac.uk

* Correspondence: zariwam@wmin.ac.uk; Tel.: +44-20-7911-5000 (ext. 65086)

† These authors contributed equally to this work.

Received: 19 June 2020; Accepted: 16 July 2020; Published: 22 July 2020



Abstract: Brain-derived neurotrophic factor (BDNF) is key for the maintenance of normal neuronal function and energy homeostasis and has been suggested to improve cognitive function, including learning and memory. Iron and the antioxidant curcumin have been shown to influence BDNF homeostasis. This 6-week, double blind, randomized, placebo-controlled study examined the effects of oral iron supplementation at low (18 mg) and high (65 mg) ferrous (FS) iron dosages, compared to a combination of these iron doses with a bioavailable formulated form of curcumin (HydroCurcTM; 500 mg) on BDNF levels in a healthy adult cohort of 155 male (26.42 years \pm 0.55) and female (25.82 years \pm 0.54) participants. Participants were randomly allocated to five different treatment groups: both iron and curcumin placebo (FS0+Plac), low dose iron and curcumin placebo (FS18+Plac), low dose iron and curcumin (FS18+Curc), high dose iron and curcumin placebo (FS65+Plac) and high dose iron and curcumin (FS65+Curc). Results showed a significant increase in BDNF over time (26%) in the FS18+Curc group ($p = 0.024$), and at end-point between FS18+Curc and FS18+Plac groups (35%, $p = 0.042$), demonstrating for the first time that the combination with curcumin, rather than iron supplementation alone, results in increased serum BDNF. The addition of curcumin to iron supplementation may therefore provide a novel approach to further enhance the benefits associated with increased BDNF levels.

Keywords: BDNF (brain-derived neurotrophic factor); curcumin; iron; ferrous sulphate; supplementation; brain function; antioxidant capability

1. Introduction

Iron is a critical micronutrient vital for oxygen transport and energy production via cellular respiration [1] as well as for the development and maintenance of normal neuronal function [2]. Assuming a mixed human diet, dietary iron intake ranges from 12 to 18 mg/day, of which 1–2 mg is absorbed into the circulation via the duodenum and proximal jejunum [3,4]. Iron intake often does not meet the body's requirements due to inadequate nutrient intake and poor bioavailability on account of the complexities of iron absorption (such as the influence of dietary iron inhibitors and enhancers), thus leading to widespread iron deficiency [5–7]. This is a worldwide public health issue as iron

deficiency has been attributed to more than 60% of anaemia cases, a condition estimated to affect around two billion people globally [8–11].

Iron deficiency can lead to impaired cognitive and physical development in children, compromise physical and cognitive performance in adults [12–15] and has been linked with fatigue [16], impaired quality of life [17] and reduced mood [14,15]. In addition to being a critical component of haemoglobin [1], iron also has a key role in electron transport during cellular respiration [18], deoxyribonucleic acid (DNA) synthesis [19] and is an important cofactor in the synthesis of neurotransmitters such as serotonin and norepinephrine [20].

Hippocampal brain-derived neurotrophic factor (BDNF) is a neurotrophic growth factor, which is suggested to be essential for normal neuronal development and cognitive function [21]. BDNF expression is associated with neurotransmitter levels and its synthesis is activated by neuronal activity and by increased cytoplasmic calcium levels via the activation of the transcription factor cyclic adenosine monophosphate (cAMP) responsive element binding protein (CREB) [22,23]. As well as being essential for maintaining the normal function of mature neurons, BDNF also regulates dendrite growth, spinal development and aids long term synaptic potentiation during neuronal development, which is associated with learning and memory formation [21]. BDNF exhibits neuroprotective effects via its role in the modulation of synaptic plasticity and function [24] and has been found to enhance explicit memory encoding, storage and retrieval of information in the hippocampal region of the brain [25]. In addition to its role in cognitive function, BDNF is thought to be a major contributor to energy homeostasis [26]. Therefore, it is thought that increased levels of BDNF could enhance cognitive capacity and potentially lead to reduced fatigue [24]. It has been shown that radiolabelled BDNF rapidly crosses the blood-brain barrier (BBB) and the efflux of unmodified BDNF to the circulation following intracerebroventricular injection has also been demonstrated in mice [27]. Further animal studies have also demonstrated the influx of BDNF from the circulation to the brain and vice versa and suggested that this may occur via a saturable transport mechanism [28,29]. Furthermore, positive correlations and parallel changes between peripheral and central BDNF levels have been shown in animal studies, indicating that circulatory BDNF measured in plasma or serum can be used as a biomarker for central BDNF levels [29–31].

Although the mechanistic pathways of iron and BDNF interaction are not yet fully elucidated, optimal iron levels are thought to be essential for BDNF homeostasis [32]. It has been suggested that when brain iron levels are low, BDNF may be down regulated as a consequence of altered neurotransmitter levels [33]. However, excessive cerebral iron levels can also reduce BDNF expression and are associated with cognitive and mental impairments [34–36]. This is claimed to be a consequence of the brain's altered redox state, as high iron levels facilitate the production of detrimental reactive oxygen species (ROS) [36,37], and the brain lacks efficient levels of antioxidant defences to counteract them [35,36]. The accumulation of ROS results in oxidative stress, damage to DNA, proteins and lipids, which can result in cell death [38,39]. Antioxidants can counteract the effects of ROS and have therefore generated interest as molecules that could potentially enhance BDNF levels [40,41].

Curcumin, a non-flavonoid polyphenol, is the most biologically active antioxidant component in the rhizomatous spice *Curcuma longa* Linnaeus (L.) or turmeric [42]. Curcumin is a pleiotropic compound with wide ranging beneficial properties including antioxidative, anti-inflammatory and neuroprotective effects [43,44]. Curcumin and its analogue curcuminoids, demethoxycurcumin (DMC) and bisdemethoxycurcumin (BDMC), can exist in two different forms depending on their environment; keto and enol [45]. In their enol form, they are capable of accepting as well as donating hydrogen and have metal chelation characteristics [46]. Curcumin is extensively studied for its ROS scavenging properties [47–50]. It acts as a free-radical chain breaker, capable of donating hydrogen to ROS due to the presence of a hydroxyl group in its structure [51]. Furthermore, studies examining the neuroprotective properties of curcumin have shown that it has potential as a treatment for neurological disorders, such as depression, bipolar disorders and neurodegenerative diseases [44,52]. Since curcumin is lipophilic in nature, it is assumed to be able to cross the BBB [53]. However, it is debated whether it

can access the brain at therapeutic concentrations as it is rapidly metabolised following ingestion [54]. The pharmacological use of curcumin has therefore been limited due to its poor bioavailability, limited bio-distribution, poor stability and short half-life [55]. Formulation science strategies have shown that nanocarrier delivery systems such as liposomes and micelles may address these limitations and enhance curcumin's therapeutic potential [44,56,57]. A recent study by Briskey and colleagues [58] demonstrated that formulation of curcumin using a novel delivery system comprising a mixture of surfactants, polar lipids and solvents (known as LipiSpense®) significantly increases the plasma concentration of curcumin in human volunteers, further highlighting the application of delivery systems to increase the bioavailability of poorly absorbed molecules.

As iron and curcumin have both been independently associated with BDNF homeostasis in animal and cellular models [33–36,52], there is scope to further investigate their co-administration. The aim of this 6-week, double blind, randomized, placebo-controlled study was to examine the effects of oral iron supplementation at low (18 mg) versus high (65 mg) ferrous iron dosages, either alone or co-administered with curcumin (500 mg) supplementation on serum BDNF levels in healthy adults. The study was designed to determine whether co-administration of ferrous sulphate and a bioavailable formulated curcumin supplement (commercially available as HydroCurc™) would amplify serum levels of BDNF.

2. Materials and Methods

2.1. Study Design

The present double blind, placebo-controlled, randomized study recruited 155 healthy participants (79 males and 76 females) for a study duration of 6 weeks. Sample size was calculated using G*Power statistical analysis software [59,60] to achieve 80% power.

Study participants were randomly allocated to one of five different treatment groups using the online service by Study Randomizer [61], via a permuted block and gender-balanced randomisation algorithm with 31 participants in each group. The 5 different treatment groups were ferrous sulphate and curcumin placebos (FS0+Plac), low ferrous sulphate (18 mg elemental iron) and curcumin placebo (FS18+Plac), low ferrous sulphate (18 mg elemental iron) and 500 mg curcumin (FS18+Curc), high ferrous sulphate (65 mg elemental iron) and curcumin placebo (FS65+Plac) and high ferrous sulphate (65 mg elemental iron) and 500 mg curcumin (FS65+Curc) (Figure 1).

Healthy adults aged between 18 and 40 years with ferritin levels in the normal physiological range were recruited into the study. Normal ferritin was defined as 15–300 µg/L for men and 15–200 µg/L for women, according to United Kingdom (UK) guidelines [62,63]. Any participants with haemoglobin values below the World Health Organization (WHO) cut-off for anaemia [64] were not enrolled in the study (<130 g/L for men and <120 g/L for women). For the purpose of sub-analysis, participants (within their allocated treatment groups), were categorised according to baseline ferritin levels; <50 µg/L was classified as 'low' and ≥50 µg/L as 'normal' ferritin values [65–69].

On the screening day, participants were assessed to ensure they complied with the inclusion/exclusion criteria by means of a comprehensive interview. Exclusion criteria included the following: any diagnosis of medical conditions or comorbidities, currently trying to conceive, pregnancy or lactating, and/or any chronic menstrual disorders or menopausal changes. Furthermore, those who had issues related to oral supplement ingestion, were on any medication or supplementation, exceeded 21 units/week of alcohol consumption, experienced any chronic gastrointestinal problems, eating disorders, psychological conditions or presented with hypo/hypertensive blood pressure, were excluded from study [70,71].

Participants signed written informed consent and ethical approval (ID: ETH1718-0907) was granted by the Faculty of Science and Technology Ethics Committee, University of Westminster, in accordance with the ethical standards of the Helsinki Declaration of 1975.




Figure 1. Study design. Participants who met the inclusion criteria during the screening day were randomly assigned to 5 different treatment groups ($n = 31/\text{group}$). There were three visit days over the study duration of 6 weeks. On the first visit day, (Baseline) body measurements, blood samples and questionnaires were collected from the participants. On the following visit days, (Mid-point and End-point) blood samples and questionnaires were collected from the participants.

2.2. Supplementation

As per Figure 1, participants were provided two different doses (high and low) of ferrous sulphate supplement, co-administered with a dose of curcumin or equivalent placebo(s) (depending on supplement group allocation). The high dose ferrous sulphate supplementation (200 mg/day, 65 mg elemental iron) is based upon the traditional first line oral iron therapy for treatment and prophylaxis of iron deficiency and iron deficiency anaemia [72,73]. The low dose ferrous sulphate supplementation (55 mg/day with 18 mg elemental iron) is based upon the recommended Daily Value (DV) of iron as per the Food and Drug Administration (FDA) [74]. Curcumin supplements were in the form of 500 mg/day of formulated curcumin (HydroCurc™, Pharmako Biotechnologies Pty Ltd. New South Wales, Australia). This formulation consisted of 85% total curcuminoids (80% curcumin, 17% DMC and 3% BDMC) entrapped in a proprietary delivery system (LipiSpense®, Pharmako Biotechnologies Pty Ltd. New South Wales, Australia) that has previously been shown to have enhanced bioavailability and deliver a higher therapeutic curcumin dose [58]. Microcrystalline cellulose served as a placebo as well as the bulking agent in the capsules of active ingredients. White-opaque hydroxypropyl methylcellulose (HPMC) capsules were used, with sizes of #1 and #00, for the ferrous sulphate and curcumin supplements, respectively. The supplements were presented in white, screw lid bottles, labelled with the related group codes. The participants were required to take one ferrous sulphate and one curcumin supplement per day with water, at least 2 h after or 1 h before food consumption; at separate times.

2.3. Physical Examination

Baseline anthropometric measurements were collected by trained research staff. Height was measured using a Seca 287 ultrasonic stadiometer (Seca GmbH & Co. KG, Hamburg, Germany). Weight and body mass index (BMI), were measured using the Seca 515 medical Body Composition Analyser (Seca GmbH & Co. KG, Hamburg, Germany). Blood pressure (BP) was measured using an Omron M6 BP monitor (Omron, Hoofddorp, The Netherlands).

2.4. Blood Collection

Participants attended blood collection appointments following an overnight fast (12 h fast). Venous blood samples were collected at baseline, mid-point (21 day) and end-point (42 day) visits from the antecubital fossa by venepuncture (using a 21G needle). Approximately 10 mL of blood was collected from each participant per procedure using Becton Dickinson (BD) Vacutainer[®] serum-separating tubes (SST) (BD, Oxford, UK). Blood in the SST was left to coagulate at room temperature for 45 min and then centrifuged (Hettich 340r, Hettich GmbH & Co. KG, Tuttlingen, Germany) for 10 min at 3857 g. Serum supernatant was aliquoted into 1.5 mL microcentrifuge tubes post-centrifugation and stored at -80°C .

2.5. Ferritin Assay

Serum ferritin samples were analysed using a Horiba ABX Pentra 400 (Horiba Ltd., Kyoto, Japan) multiparametric medical bench top chemistry analyser, compliant with the National Committee for Clinical Laboratory Standards (NCCLS) [75]. Ferritin values were determined by latex-enhanced immunoturbidimetric assay, in accordance with the manufacturer's protocol and as previously described by Simó et al. [76].

2.6. BDNF Assay

Serum was assayed for BDNF levels using the Biosensis Mature BDNF Rapid[™] enzyme-linked immunosorbent assay (ELISA) Kit (ATI Atlas, Chichester, UK) following the manufacturer's protocol, using a dilution factor of 1:100. Pre-coated microplates were incubated with 100 μL of diluted BDNF standards, quality control (QC) samples, serum samples (1:100) or blanks (sample diluent only) for 45 min on a plate shaker (140 rpm), at room temperature (RT). Plates were then washed five times with wash buffer (200 μL per well). After the addition of 100 μL detection antibody per well, the plates were incubated on the plate shaker (140 rpm at RT) for 30 min. Following five more washes, 100 μL of 1 \times streptavidin-HRP conjugate was added to each well. The plates were incubated for a further 30 min at 140 rpm (RT). The plates were then washed 5 times and 100 μL of 3,3',5,5'-Tetramethylbenzidine (TMB) was added to each well and incubated at RT for 9 min in the dark before the addition of 100 μL of stop solution into each well. The absorbance was read with a SPECTROstar[®] Nano microplate reader (BMG Labtech GmbH, Ortenberg, Germany) at 450 nm (within 5 min).

2.7. Study Compliance

Compliance with the study protocol (including supplementation) and adverse effects were checked during the mid-point (week 3) and at the end-point (week 6) of each study period and any deviation from the study protocol was noted and assessed. Adherence to daily supplementation of the active or placebo capsules was $\geq 80\%$ at all times. Data from all participants who successfully completed the study period was utilised for further analysis.

2.8. Statistical Analysis

Values are expressed as mean \pm Standard Error of Mean (SEM). The BDNF assay results were statistically analysed using a two-way, repeated measures analysis of variance (ANOVA) or mixed effects model (where missing values were present). Post-hoc tests (Sidak's and Tukey's) were carried out to assess differences between and within treatment groups (PRISM software package, Version 8, Graphpad Software Inc., San Diego, CA, USA).

3. Results

Of the 155 participants recruited, 150 completed all study visits. Two participants withdrew from the FS0+Plac groups, one from nausea after the baseline visit and one from loss of interest in the study after the mid-point visit. One participant also withdrew from the FS65+Curc group after the baseline

visit due to loss of interest and another withdrew due to gastric distress. A participant in the FS18+Plac group was excluded from the study due to incomplete blood sampling at the mid-point (Figure S1).

The mean age of participants was 26.12 years (± 0.39). There was no significant difference in mean age between the five treatment groups. There were also no significant differences observed in anthropometric measurements of participants (Table 1). The study population was of mixed ethnicity, representative of the population at the site of recruitment (London, UK).

Table 1. Participant age, weight, height, body mass index (BMI) and body fat percentages (mean \pm SEM).

Variable	FS0+Plac	FS18+Plac	FS18+Curc	FS65+Plac	FS65+Curc
Age (yrs)	26.29 \pm 0.84	25.84 \pm 0.93	24.48 \pm 0.82	27.23 \pm 0.83	26.77 \pm 0.87
Weight (kg)	70.79 \pm 2.37	72.45 \pm 3.13	66.17 \pm 2.45	70.21 \pm 3.54	67.70 \pm 2.13
Height (m)	1.72 \pm 0.02	1.71 \pm 0.01	1.68 \pm 0.02	1.70 \pm 0.02	1.72 \pm 0.02
BMI (m/kg ²)	23.89 \pm 0.58	24.51 \pm 0.85	23.32 \pm 0.64	24.11 \pm 1.00	22.83 \pm 0.55
Body fat (%)	25.27 \pm 1.71	24.96 \pm 1.67	24.32 \pm 1.76	24.97 \pm 1.54	23.39 \pm 1.39

At baseline, no significant differences were observed in mean ferritin levels across the treatment groups (Table 2). However, there was a significant difference observed in baseline BDNF levels between the FS18+Plac (37.16 ng/mL) and FS18+Curc (30.28 ng/mL) groups, with the mean BDNF being (22.7% $p = 0.049$) higher in the FS18+Plac group compared to the FS18+Curc group (Table 2). No significant differences in baseline BDNF were observed when comparing any of the other groups (Table 2).

Table 2. Participant baseline ferritin and brain-derived neurotrophic factor (BDNF) levels expressed as mean \pm SEM.

Variable	FS0+Plac	FS18+Plac	FS18+Curc	FS65+Plac	FS65+Curc
Ferritin(μ g/L)	58.71 \pm 9.37	68.58 \pm 10.30	52.36 \pm 7.74	55.14 \pm 8.18	61.46 \pm 8.47
BDNF (ng/mL)	35.18 \pm 2.10	37.16 \pm 1.88	30.28 \pm 1.54	31.59 \pm 1.35	30.85 \pm 8.47

When evaluating the effect of treatment group on serum BDNF levels after 21 day (mid-point) and 42 day (end-point) supplementation, significant differences were observed between the different treatment groups ($F(4, 144) = 2.746, p = 0.031$) and the two time points ($F(1, 142) = 11.36, p = 0.001$). A significant increase of 26.34% in BDNF levels from mid-point to end-point was observed in participants taking FS18+Curc ($p = 0.024$) (Figure 2). At the end-point, there was also a significant difference observed in BDNF levels between the FS18+Curc and FS18+Plac groups ($p = 0.042$), with the FS18+Curc treatment resulting in a 34.94% higher concentration of BDNF than FS18+Plac (Figure 2). A similar trend of increased BDNF was observed at the end-point when comparing the FS65+Curc treatment with FS65+Plac, however this was not significant (Figure 2). There were no other significant differences in BDNF levels between or within treatment groups (Figure 2).

When participants were sub-grouped according to low ferritin ($<50 \mu$ g/L) and normal ferritin ($\geq 50 \mu$ g/L) values, a significant increase in BDNF from mid-point to end-point was observed in the cohort with low ferritin who were supplemented with FS18+Curc ($p = 0.019$) (Figure 3A). Although no significant difference in BDNF was observed between the FS18+Curc and FS18+Plac group for participants with low baseline ferritin, the FS18+Curc group had significantly higher BDNF at the end-point compared to the FS0+Plac group, (increased by 53.78%, $p = 0.028$) (Figure 3A). No significant differences in BDNF levels were observed between treatment groups, at either time point, in participants with normal ferritin levels (Figure 3B).




Figure 2. BDNF levels are expressed as percentage difference compared to baseline levels (mean ± SEM): FS0+Plac [Mid-point *n* = 29; End-point *n* = 28], FS18+Plac [Mid-point *n* = 30; End-point *n* = 29], FS18 + Curc [Mid-point *n* = 31; End-point *n* = 31], FS65+Plac [Mid-point *n* = 30; End-point *n* = 30] and FS65 + Curc [Mid-point *n* = 29; End-point *n* = 29]). Samples were collected and analysed at mid-point (day 21) and end-point (day 42). * represents significance values when comparing each condition and time points within the same condition. (* *p* < 0.05).

No significant differences were observed in baseline ferritin levels between the low and normal ferritin groups (Tables 3 and 4). In the low ferritin sub-group (Table 3), significant differences were observed between baseline, mid-point and end-point time points ($F(2, 158) = 27.81, p < 0.0001$). A significant increase of 35.17% in ferritin levels was observed at end-point compared to baseline in the FS18+Curc group ($p = 0.0013$) (Table 3). A significant increase of 58.75% in ferritin levels was also observed at end-point compared to baseline in the FS65+Curc group ($p = 0.0002$) (Table 3).

Furthermore, there were significant increases from baseline for the FS65+Plac group at mid-point (43.6%) and end-point (68.34%) ($p = 0.0014$ and $p < 0.0001$, respectively) (Table 3). In the normal ferritin sub-group, there was no significant effect observed in ferritin levels over time or between groups in relation to any supplementation (Table 4).

Table 3. Mean ferritin (µg/L) values (low ferritin sub-group) per treatment group and timepoints (mean ± SEM). * represents significance values when comparing mid-point or end-point to baseline within the same condition (** *p* < 0.01; *** *p* < 0.001; **** *p* < 0.0001).

Timepoint	FS0+Plac	FS18+Plac	FS18+Curc	FS65+Plac	FS65+Curc
Baseline	25.08 ± 1.73	25.29 ± 2.81	30.25 ± 2.52	25.55 ± 1.78	25.04 ± 2.72
Mid-point	28.23 ± 2.96	34.59 ± 4.24	36.45 ± 2.92	36.69 ± 3.34 **	32.10 ± 4.04
End-point	31.39 ± 5.12	33.31 ± 3.39	40.89 ± 4.99 **	43.01 ± 4.01 ****	39.75 ± 6.19 ***

Table 4. Mean ferritin (µg/L) values (normal ferritin sub-group) per treatment group and timepoints (mean ± SEM).

Timepoint	FS0+Plac	FS18+Plac	FS18+Curc	FS65+Plac	FS65+Curc
Baseline	91.38 ± 12.94	109.17 ± 13.30	98.79 ± 15.37	94.54 ± 9.94	100.31 ± 10.01
Mid-point	93.37 ± 13.75	100.10 ± 16.85	102.23 ± 13.56	101.50 ± 11.75	101.63 ± 11.45
End-point	87.88 ± 11.45	100.21 ± 13.27	102.32 ± 16.18	98.37 ± 15.49	106.62 ± 12.29




Figure 3. (A). Mean \pm SEM, % difference in BDNF for participants with low ferritin at baseline ($<50 \mu\text{g/L}$): FS0+Plac [Mid-point $n = 16$; End-point: $n = 15$]; FS18+Plac [Mid-point $n = 14$; End-point $n = 13$]; FS18+Curc [Mid-point $n = 21$; End-point $n = 21$]; FS65+Plac [Mid-point $n = 19$; End-point $n = 19$] and FS65+Curc [Mid-point $n = 14$; End-point $n = 14$]. Samples were collected and analysed at mid-point (day 21) and end-point (day 42). (B). Mean \pm SEM, % difference in BDNF for participants with normal ferritin ($\geq 50 \mu\text{g/L}$): FS0+Plac [Mid and End-point $n = 15$]; FS18+Plac [Mid and End-point $n = 16$]; FS18+Curc [Mid and End-point $n = 10$]; FS65+Plac [Mid and End-point $n = 11$] and FS65+Curc [Mid and End-point $n = 15$]. * represents significance values when comparing each condition and time points within the same condition. (* $p < 0.05$).

4. Discussion

Research has shown BDNF to have a key role in the maintenance of normal neuronal function [21], energy homeostasis [26] and, in animal studies, it has been demonstrated to enhance declarative memory in the developed nervous system [25]. These findings indicate that increased BDNF levels could help to improve cognitive function. Iron supplementation in mice [32] and formulated curcumin supplementation in humans have been shown to independently increase BDNF levels [52], indicating that the combination of these two supplements could possibly express an additive effect on serum BDNF levels. The current study therefore aimed to examine the effects of iron supplement dose in the presence or absence of formulated curcumin on BDNF levels in a healthy adult population aged 18 to 40 years.

Study compliance was high with approximately 3% of initial recruits dropping out or being excluded. Although gastrointestinal effects (such as nausea or delayed bowel movements) were reported by some participants, such incidences were usually mild or transient. High compliance and mild, transient incidences seen in the current study, could be attributed to the dosages of ferrous

sulphate being within ranges that are unlikely to cause a high degree of gastric distress, which often results in the poor compliance typically seen in oral iron supplementation [77].

All participants had ferritin values in the normal range for the UK (15–300 µg/L—men and 15–200 µg/L—women) [63]. The 2011 WHO guidelines state that serum ferritin less than 15 µg/L is a specific indicator for iron deficiency [78]. Globally, there is ongoing debate calling for an increase in the normal ferritin cut-off value. Numerous authors [79–81] recommend a serum ferritin cut-off of 30 µg/L, as it has been shown to be indicative of insufficient iron stores. Iron therapy may therefore be appropriate in such individuals (ferritin <30 µg/L) to support erythropoiesis as well as normal physical performance and cognitive health [80]. It has also been argued that ferritin below 50 µg/L may indicate reduced bone marrow iron stores or latent iron deficiency [81]. Furthermore, a reduction in fatigue has been reported following iron supplementation in people with ferritin below 50 µg/L [66,68], suggesting the potential therapeutic value of iron supplementation within this ferritin range.

Similar to Verdon et al. [66] and Vaucher et al. [68], whilst taking into account the recent work from Soppi [81], data from the current study were sub-grouped into ‘low’ (<50 µg/L) ferritin and ‘normal’ (≥50 µg/L) ferritin participants (Figure 3A,B, respectively). Participants with serum ferritin below 50 µg/L showed the same trend (increased levels) as the overall data for serum BDNF, with the FS18+Curc treatment leading to increased BDNF from mid-point to end-point (Figure 3A). However, no significant differences were recorded for participants with normal ferritin levels. This suggests that the addition of curcumin to 18 mg iron supplementation, in particular, may be most effective at enhancing serum BDNF levels in individuals with low ferritin levels. Notably, in participants who had ‘low’ ferritin at baseline, all groups containing iron supplementation showed significant increases in ferritin values, apart from the FS18+Plac group. This indicates that the addition of curcumin alongside low dose iron supplementation may contribute to enhanced ferritin formation, which may indicate enhanced intestinal iron uptake over time. Iron status (ferritin status) can inversely influence the body’s iron absorption uptake in order to maintain homeostasis [82,83], and this may explain the lack of effect seen in participants who had ‘normal’ baseline ferritin values (Table 4). Together, these results suggest that curcumin may enhance the effects of low dose iron supplementation, in particular for those individuals with iron deficiency [77,83,84].

As iron is an essential component for mono-oxygenase enzyme production as well as a key co-factor in serotonin synthesis [20], it may be a key component of BDNF synthesis since neuroamines including norepinephrine can raise cAMP levels, leading to enhanced BDNF synthesis via increased phosphorylation of CREB [32]. Therefore, the general lack of change in BDNF levels observed overall and with participants following iron-only treatment may have occurred because iron is utilised for several biological processes following absorption, such as cellular respiration [19], and thus brain iron levels may not have been increased sufficiently to enhance neurotransmitter synthesis and BDNF expression. Compared to conventional curcumin, Briskey et al. [58] showed that a formulated curcumin form resulted in significantly greater increases in total plasma curcuminoid concentration in a human volunteer study. Furthermore, formulated curcumin has been shown to be more likely to cross the blood–brain barrier [44,54,85] and express neuroprotective effects due to improved bioavailability and targeted delivery, in both cellular and murine studies [44,85,86]. This could therefore explain how the addition of curcumin to iron supplementation consistently resulted in increased BDNF levels in the current study. There are various mechanistic pathways via which curcumin expresses its neuroprotective properties. The increase in serum BDNF observed following FS18+Curc treatment is supportive of previous studies, which have shown curcumin to increase BDNF synthesis by increasing cAMP levels and activating extracellular signal-regulated kinases (ERKs) and p38 mitogen-activated protein kinase (p38 MAPK), which are known to induce nuclear translocation of transcription factors, such as ETS-like protein-1 (Elk-1), activator protein 1 (AP-1) and nuclear factor kappa B (NF_κB) [52,87]. However, as there was an absence of a curcumin-only group in the current study, it cannot be confirmed that the observed increase in serum BDNF is solely a result of curcumin and not the result of synergic effects of the co-administration of iron and formulated curcumin supplementation.

The current study also demonstrated an increase in serum BDNF levels between mid- and end-point in the high dose iron group (FS65+Curc), with an observable difference between FS65+Curc and FS65+Plac groups at the end-point. However, these results were not significant. This observable but non-significant change could be due to reduced fractional iron absorption (FIA); in other words, the proportion of the dose absorbed. According to Moretti et al. [88], iron doses above 60 mg, when taken on consecutive rather than alternative days, resulted in reduced FIA. This is understood to be mediated by significantly increased levels of serum hepcidin, which are observed following 24 h supplementation with ≥ 60 mg elemental iron [88]. Hepcidin is the key regulatory protein of systemic iron homeostasis and protects against iron overload [89]. It is synthesised mainly by the liver and regulated by iron, erythropoietic activity, hypoxia and inflammation [90]. Hepcidin regulates iron homeostasis by binding to ferroportin, the transmembrane protein that mediates iron efflux from the cells, initiating its internalisation and degradation. Consequently, the iron-efflux activity of ferroportin is down regulated, leading to a reduced dose of iron being transported to the systemic circulation from the gut enterocytes [90,91]. Stoffel et al. [92] showed that FIA was 34% lower following 14 consecutive days compared to 28 alternative days dosing with oral ferrous sulphate (60 mg elemental iron supplementation) in iron-depleted women (median age 22 years) without anaemia.

Alternatively, reduced iron absorption following consecutive days of high iron supplementation could be explained by the mucosal block theory [93]. The enterocytes uptake iron from the gut lumen via divalent metal transporter 1 (DMT1) in the ferrous (Fe^{2+}) form. This iron is then converted into ferric (Fe^{3+}) form and can be stored by binding to apoferritin, forming ferritin, (the temporary storage form of iron) or can be transported to the systemic circulation via ferroportin after being reduced back to its Fe^{2+} form [94]. It is assumed that enterocytes exposed to large doses of iron would store iron in ferritin form, when the body's iron requirement is reduced. Furthermore, with the reduced availability of apoferritin, the enterocytes' iron absorption would consequently reduce until the cells are replaced in 5 to 6 days [95] with their iron content being lost in the faeces. Nonetheless, the accuracy of mucosal block theory has been debated in recent studies [96–98]. However, in either case, reduced FIA or mucosal block could lead to the accumulation of large amounts of unabsorbed iron, which may cause gut inflammation [77], increase the production of free radicals in the mucosa (the innermost layer of the gut) [99,100] and may alter the gut microbiota by increasing pathogenic microorganisms whilst reducing commensal microflora [101,102]. Commensal microorganisms such as *Escherichia coli*, *Blautia sp.* (MRG-PMF1) or *Lactobacillus acidophilus* have been shown to be biologically active in curcumin metabolism in the gut, producing curcumin derivatives with increased biological activity and improved pharmacokinetics related to absorption compared to native curcumin. Thus, microbiota may enhance the neuroprotective effects of curcumin [103]. However, the enhancing role of curcumin on gut microbiota may be diminished when compromised by high iron accumulation in the gut [104]. Additionally, the increased levels of free radicals in the gut following high iron absorption may result in the free radical scavenging properties of curcumin being utilised locally [48]. This could explain why the combination of curcumin with low iron supplementation (FS18+Curc) generally resulted in higher end-point serum BDNF levels than with high iron plus curcumin (FS65+Curc) treatment.

The normal range of serum BDNF levels varies according to numerous factors such as age, ethnicity, study location and sample processing methods [105–107]. However, previous research indicates that normal serum BDNF levels may fall within the range of 8 ng/mL and 46 ng/mL [108,109]. The data from the current study suggests that supplementation with 18 mg iron and 500 mg curcumin over 42 days elevates BDNF levels towards the higher end of this normal range, given the average end-point value of 39.17 ng/mL within this group (Table S1). However, a recent Swiss study of healthy adults reported a wider range of serum BDNF levels, between 15.83 ng/mL and 79.77 ng/mL [107], which highlights the difficulty in precisely establishing normal serum BDNF levels within a healthy cohort. Nevertheless, the relationship between BDNF and cognitive function is well established [110–113] and low serum BDNF levels have been attributed to cognitive impairment in the elderly [114]. Therefore, the evident increase in serum BDNF following low iron and curcumin supplementation in this cohort may be

expected to be associated with probable cognitive benefits. However, in the future, it would be of value to also directly assess cognitive performance in relation to serum BDNF following iron and curcumin supplementation at varying timeframes, as this could help identify the specific levels of serum BDNF that correlate with the enhancement of cognitive function.

In summary, the current study demonstrates for the first time that the co-administration of a bioavailable formulated curcumin supplement with ferrous sulphate containing 18 mg elemental iron for 42 days results in increased serum BDNF levels. The addition of curcumin may therefore provide a novel approach to iron supplementation and possibly enhance the iron-associated cognitive benefits linked to increased serum BDNF levels.

Supplementary Materials: The following are available online at <http://www.mdpi.com/2076-3921/9/8/645/s1>, Figure S1: shows the study compliance after 155 participants were enrolled and randomised equally into 5 treatment groups: FS0+Plac (full placebo, placebos for both iron and curcumin), FS18+Plac (18 mg elemental iron and placebo for curcumin), FS18+Curc (18 mg elemental iron and 500 mg curcumin), FS65+Plac (65 mg elemental iron and placebo for curcumin) and FS65+Curc (65 mg elemental iron and 500 mg curcumin). Table S1: Mean BDNF (ng/mL) values per treatment group/timepoint (mean \pm SEM).

Author Contributions: Conceptualisation and methodology: M.G.Z., D.R., G.B. and O.F. Supervision: M.G.Z., D.R. and G.B. Investigation and data curation: O.F. and H.T.L. Data analysis: O.F., H.T.L., L.M. and G.B. Writing—original draft preparation: H.T.L., O.F., L.M. and M.G.Z. Writing—review and editing: M.G.Z., D.R., G.B. and L.M. All authors have read and agreed to the published version of the manuscript.

Funding: The study was supported by Knowledge Exchange funding from Gencor Pacific Ltd (UT722720).

Acknowledgments: We gratefully acknowledge the time and dedication of all participants who participated in this study. We would like to acknowledge all colleagues from The University of Westminster who contributed support and expertise to this study. We thank Lina Temouri for assistance with administrative duties, phlebotomy and laboratory work, Chrystalla Ferrier for assisting with the operation of the Ilab Aries instrument and Yvoni Kyriakidou for assistance with phlebotomy. We thank Sonita Chea, Hashum Sum and Ruchita Venkatesh for their laboratory and administration assistance and Gloria Koncz for her administration assistance, and Mariana Peixoto for her logistical support to this project. The authors would also like to thank the Research Interns and Visiting Researchers who assisted over the course of this project.

Conflicts of Interest: The funder was involved in the study design process but had no involvement in the data collection or analysis. The authors declare no other conflict of interest.

References

1. Ganz, T.; Nemeth, E. Iron homeostasis in host defence and inflammation. *Nat. Rev. Immunol.* **2015**, *15*, 500–510. [[CrossRef](#)] [[PubMed](#)]
2. Beard, J.L.; Connor, J.R. Iron status and neural functioning. *Annu. Rev. Nutr.* **2003**, *23*, 41–58. [[CrossRef](#)] [[PubMed](#)]
3. Zariwala, M.; Elsaid, N.; Jackson, T.L.; López, F.C.; Farnaud, S.; Somavarapu, S.; Renshaw, D. A novel approach to oral iron delivery using ferrous sulphate loaded solid lipid nanoparticles. *Int. J. Pharm.* **2013**, *456*, 400–407. [[CrossRef](#)] [[PubMed](#)]
4. Dev, S.; Babitt, J.L. Overview of iron metabolism in health and disease. *Hemodial. Int.* **2017**, *21*, S6–S20. [[CrossRef](#)]
5. Zimmermann, M.; Hurrell, R. Nutritional iron deficiency. *Lancet* **2007**, *370*, 511–520. [[CrossRef](#)]
6. Hurrell, R.F.; Egli, I. Iron bioavailability and dietary reference values. *Am. J. Clin. Nutr.* **2010**, *91*, 1461S–1467S. [[CrossRef](#)]
7. Paesano, R.; Natalizi, T.; Berlutti, F.; Valenti, P. Body iron delocalization: The serious drawback in iron disorders in both developing and developed countries. *Pathog. Glob. Health* **2012**, *106*, 200–216. [[CrossRef](#)]
8. McLean, E.; Cogswell, M.; Egli, I.; Wojdyla, D.; De Benoist, B. Worldwide prevalence of anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993–2005. *Public Health Nutr.* **2009**, *12*, 444–454. [[CrossRef](#)]
9. de Benoist, B.; McLean, E.; Egli, I.; Cogswell, M. *World Health Organization Worldwide Prevalence of Anaemia 1993–2005: WHO Global Database on Anaemia*; WHO Press, World Health Organization: Geneva, Switzerland, 2008.
10. Kassebaum, N.J. The Global Burden of Anemia. *Hematol. Oncol. Clin.* **2016**, *30*, 247–308. [[CrossRef](#)]
11. Williams, A.M.; Addo, O.Y.; Grosse, S.D.; Kassebaum, N.J.; Rankin, Z.; Ballesteros, K.E.; Olsen, H.E.; Sharma, A.J.; Jefferds, M.E.; Mei, Z. Data needed to respond appropriately to anemia when it is a public health problem. *Ann. N. Y. Acad. Sci.* **2019**, *1450*, 268–280. [[CrossRef](#)]

12. Haas, J.D.; Brownlie, T. Iron deficiency and reduced work capacity: A critical review of the research to determine a causal relationship. *J. Nutr.* **2001**, *131*, 676S–690S. [[CrossRef](#)] [[PubMed](#)]
13. Brownlie, T.; Utermohlen, V.; Hinton, P.; Giordano, C.; Haas, J.D. Marginal iron deficiency without anemia impairs aerobic adaptation among previously untrained women. *Am. J. Clin. Nutr.* **2002**, *75*, 734–742. [[CrossRef](#)] [[PubMed](#)]
14. Jáuregui-Lobera, I. Iron deficiency and cognitive functions. *Neuropsychiatr. Dis. Treat.* **2014**, *10*, 2087–2095. [[CrossRef](#)]
15. Lopez, A.; Cacoub, P.; Macdougall, I.C.; Peyrin-Biroulet, L. Iron deficiency anaemia. *Lancet* **2016**, *387*, 907–916. [[CrossRef](#)]
16. Patterson, A.J.; Brown, W.J.; Powers, J.R.; Roberts, D.C. Iron deficiency, general health and fatigue: Results from the Australian Longitudinal Study on Women’s Health. *Qual. Life Res.* **2000**, *9*, 491–497. [[CrossRef](#)] [[PubMed](#)]
17. Enjuanes, C.; Klip, I.T.; Bruguera, J.; Cladellas, M.; Ponikowski, P.; Banasiak, W.; Van Veldhuisen, D.J.; Van Der Meer, P.; Jankowska, E.A.; Comin-Colet, J. Iron deficiency and health-related quality of life in chronic heart failure: Results from a multicenter European study. *Int. J. Cardiol.* **2014**, *174*, 268–275. [[CrossRef](#)] [[PubMed](#)]
18. Pino, J.M.V.; Da Luz, M.H.M.; Antunes, H.K.M.; Giampá, S.Q.D.C.; Martins, V.R.; Lee, K.S. Iron-Restricted Diet Affects Brain Ferritin Levels, Dopamine Metabolism and Cellular Prion Protein in a Region-Specific Manner. *Front. Mol. Neurosci.* **2017**, *10*, 145. [[CrossRef](#)]
19. Abbaspour, N.; Hurrell, R.; Kelishadi, R. Review on iron and its importance for human health. *J. Res. Med. Sci.* **2014**, *19*, 164–174.
20. Chen, M.J.; Nguyen, T.; Pike, C.J.; Russo-Neustadt, A. Norepinephrine induces BDNF and activates the PI-3K and MAPK cascades in embryonic hippocampal neurons. *Cell. Signal.* **2007**, *19*, 114–128. [[CrossRef](#)]
21. Gonzalez, A.; Moya-Alvarado, G.; Gonzalez-Billaut, C.; Bronfman, F.C. Cellular and molecular mechanisms regulating neuronal growth by brain-derived neurotrophic factor. *Cytoskeleton* **2016**, *73*, 612–628. [[CrossRef](#)]
22. Tao, X.; Finkbeiner, S.; Arnold, D.B.; Shaywitz, A.J.; Greenberg, M.E. Ca²⁺ Influx Regulates BDNF Transcription by a CREB Family Transcription Factor-Dependent Mechanism. *Neuron* **1998**, *20*, 709–726. [[CrossRef](#)]
23. Zheng, F.; Zhou, X.; Luo, Y.; Xiao, H.; Wayman, G.; Wang, H. Regulation of Brain-Derived Neurotrophic Factor Exon IV Transcription through Calcium Responsive Elements in Cortical Neurons. *PLoS ONE* **2011**, *6*, e28441. [[CrossRef](#)] [[PubMed](#)]
24. Bathina, S.; Das, U.N. Brain-derived neurotrophic factor and its clinical implications. *Arch. Med. Sci.* **2015**, *11*, 1164–1178. [[CrossRef](#)] [[PubMed](#)]
25. Bekinschtein, P.; Cammarota, M.; Kathe, C.; Slipczuk, L.; Rossato, J.I.; Goldin, A.; Izquierdo, I.; Medina, J.H. BDNF is essential to promote persistence of long-term memory storage. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 2711–2716. [[CrossRef](#)] [[PubMed](#)]
26. Bothwell, M. Functional interactions of neurotrophins and neurotrophin receptors. *Annu. Rev. Neurosci.* **1995**, *18*, 223–253. [[CrossRef](#)] [[PubMed](#)]
27. Pan, W.; Banks, W.A.; Fasold, M.B.; Bluth, J.; Kastin, A.J. Transport of brain-derived neurotrophic factor across the blood-brain barrier. *Neuropharmacology* **1998**, *37*, 1553–1561. [[CrossRef](#)]
28. Poduslo, J.F.; Curran, G.L. Permeability at the blood-brain and blood-nerve barriers of the neurotrophic factors: NGF, CNTF, NT-3, BDNF. *Mol. Brain Res.* **1996**, *36*, 280–286. [[CrossRef](#)]
29. Sartorius, A.; Hellweg, R.; Litzke, J.; Vogt, M.A.; Dormann, C.; Vollmayr, B.; Danker-Hopfe, H.; Gass, P. Correlations and Discrepancies between Serum and Brain Tissue Levels of Neurotrophins after Electroconvulsive Treatment in Rats. *Pharmacopsychiatry* **2009**, *42*, 270–276. [[CrossRef](#)]
30. Karege, F.; Schwald, M.; Cisse, M. Postnatal developmental profile of brain-derived neurotrophic factor in rat brain and platelets. *Neurosci. Lett.* **2002**, *328*, 261–264. [[CrossRef](#)]
31. Klein, A.B.; Williamson, R.; Santini, M.A.; Clemmensen, C.; Ettrup, A.; Rios, M.; Knudsen, G.M.; Aznar, S. Blood BDNF concentrations reflect brain-tissue BDNF levels across species. *Int. J. Neuropsychopharmacol.* **2011**, *14*, 347–353. [[CrossRef](#)]
32. Mehrpouya, S.; Nahavandi, A.; Khojasteh, F.; Soleimani, M.; Ahmadi, M.; Barati, M. Iron administration prevents BDNF decrease and depressive-like behavior following chronic stress. *Brain Res.* **2015**, *1596*, 79–87. [[CrossRef](#)] [[PubMed](#)]

33. Texel, S.J.; Camandola, S.; Ladenheim, B.; Rothman, S.M.; Mughal, M.R.; Unger, E.L.; Cadet, J.L.; Mattson, M.P. Ceruloplasmin deficiency results in an anxiety phenotype involving deficits in hippocampal iron, serotonin, and BDNF. *J. Neurochem.* **2012**, *120*, 125–134. [[CrossRef](#)] [[PubMed](#)]
34. Lozoff, B.; Jimenez, E.; Wolf, A.W. Long-Term Developmental Outcome of Infants with Iron Deficiency. *N. Engl. J. Med.* **1991**, *325*, 687–694. [[CrossRef](#)] [[PubMed](#)]
35. Sian-Hülsmann, J.; Mandel, S.; Youdim, M.B.H.; Riederer, P. The relevance of iron in the pathogenesis of Parkinson's disease. *J. Neurochem.* **2011**, *118*, 939–957. [[CrossRef](#)] [[PubMed](#)]
36. Radak, Z.; Suzuki, K.; Higuchi, M.; Balogh, L.; Boldogh, I.; Koltai, E. Physical exercise, reactive oxygen species and neuroprotection. *Free. Radic. Biol. Med.* **2016**, *98*, 187–196. [[CrossRef](#)] [[PubMed](#)]
37. Hwang, O. Role of oxidative stress in Parkinson's disease. *Exp. Neurobiol.* **2013**, *22*, 11–17. [[CrossRef](#)]
38. Ray, P.D.; Huang, B.-W.; Tsuji, Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell. Signal.* **2012**, *24*, 981–990. [[CrossRef](#)]
39. Jiang, H.; Wang, J.; Rogers, J.; Xie, J. Brain Iron Metabolism Dysfunction in Parkinson's Disease. *Mol. Neurobiol.* **2017**, *54*, 3078–3101. [[CrossRef](#)]
40. Fahnestock, M.; Marchese, M.; Head, E.; Pop, V.; Michalski, B.; Milgram, W.N.; Cotman, C.W. BDNF increases with behavioral enrichment and an antioxidant diet in the aged dog. *Neurobiol. Aging* **2012**, *33*, 546–554. [[CrossRef](#)]
41. Assunção, M.; Santos-Marques, M.J.; Carvalho, F.; Andrade, J.P. Green tea averts age-dependent decline of hippocampal signaling systems related to antioxidant defenses and survival. *Free Radic. Biol. Med.* **2010**, *48*, 831–838. [[CrossRef](#)]
42. Kulkarni, S.K.; Dhir, A.; Akula, K.K. Potentials of Curcumin as an Antidepressant. Available online: <https://www.hindawi.com/journals/tswj/2009/624894/> (accessed on 8 April 2020).
43. Lopresti, A.L.; Hood, S.; Drummond, P.D. Multiple antidepressant potential modes of action of curcumin: A review of its anti-inflammatory, monoaminergic, antioxidant, immune-modulating and neuroprotective effects. *J. Psychopharmacol.* **2012**, *26*, 1512–1524. [[CrossRef](#)] [[PubMed](#)]
44. Mursaleen, L.; Somavarapu, S.; Zariwala, M.G. Deferoxamine and Curcumin Loaded Nanocarriers Protect Against Rotenone-Induced Neurotoxicity. *J. Parkinsons Dis.* **2020**, *10*, 99–111. [[CrossRef](#)] [[PubMed](#)]
45. Bich, V.T.; Thuy, N.T.; Binh, N.T.; Huong, N.T.M.; Yen, P.N.D.; Luong, T.T. *Structural and Spectral Properties of Curcumin and Metal-Curcumin Complex Derived from Turmeric (Curcuma Longa)*; Springer: Berlin/Heidelberg, Germany, 2009; pp. 271–278.
46. Ferrari, E.; Benassi, R.; Sacchi, S.; Pignedoli, F.; Asti, M.; Saladini, M. Curcumin derivatives as metal-chelating agents with potential multifunctional activity for pharmaceutical applications. *J. Inorg. Biochem.* **2014**, *139*, 38–48. [[CrossRef](#)]
47. Tennesen, H.H.; Greenhill, J.V. Studies on curcumin and curcuminoids. XXII: Curcumin as a reducing agent and as a radical scavenger. *Int. J. Pharm.* **1992**, *87*, 79–87. [[CrossRef](#)]
48. Reddy, A.C.P.; Lokesh, B.R. Studies on the inhibitory effects of curcumin and eugenol on the formation of reactive oxygen species and the oxidation of ferrous iron. *Mol. Cell. Biochem.* **1994**, *137*, 1–8. [[CrossRef](#)] [[PubMed](#)]
49. Borsari, M.; Ferrari, E.; Grandi, R.; Saladini, M. Curcuminoids as potential new iron-chelating agents: Spectroscopic, polarographic and potentiometric study on their Fe (III) complexing ability. *Inorg. Chim. Acta* **2002**, *328*, 61–68. [[CrossRef](#)]
50. Bernabé-Pineda, M.; Ramírez-Silva, M.T.; Romero-Romo, M.; González-Vergara, E.; Rojas-Hernández, A. Determination of acidity constants of curcumin in aqueous solution and apparent rate constant of its decomposition. *Spectrochim. Acta Part Mol. Biomol. Spectrosc.* **2004**, *60*, 1091–1097. [[CrossRef](#)]
51. Salem, M.; Rohani, S.; Gillies, E.R. Curcumin, a promising anti-cancer therapeutic: A review of its chemical properties, bioactivity and approaches to cancer cell delivery. *RSC Adv.* **2014**, *4*, 10815–10829. [[CrossRef](#)]
52. Sarraf, P.; Parohan, M.; Javanbakht, M.H.; Ranji-Burachaloo, S.; Djalali, M. Short-term curcumin supplementation enhances serum brain-derived neurotrophic factor in adult men and women: A systematic review and dose-response meta-analysis of randomized controlled trials. *Nutr. Res.* **2019**, *69*, 1–8. [[CrossRef](#)]
53. Rossi, L.; Mazzitelli, S.; Arciello, M.; Capo, C.R.; Rotilio, G. Benefits from Dietary Polyphenols for Brain Aging and Alzheimer's Disease. *Neurochem. Res.* **2008**, *33*, 2390–2400. [[CrossRef](#)]
54. Tsai, Y.-M.; Chien, C.-F.; Lin, L.-C.; Tsai, T.-H. Curcumin and its nano-formulation: The kinetics of tissue distribution and blood–brain barrier penetration. *Int. J. Pharm.* **2011**, *416*, 331–338. [[CrossRef](#)] [[PubMed](#)]

55. Prasad, S.; Tyagi, A.K.; Aggarwal, B.B. Recent Developments in Delivery, Bioavailability, Absorption and Metabolism of Curcumin: The Golden Pigment from Golden Spice. *Cancer Res. Treat.* **2014**, *46*, 2. [[CrossRef](#)] [[PubMed](#)]
56. Zhongfa, L.; Chiu, M.; Wang, J.; Chen, W.; Yen, W.; Fan-Havard, P.; Yee, L.D.; Chan, K.K. Enhancement of curcumin oral absorption and pharmacokinetics of curcuminoids and curcumin metabolites in mice. *Cancer Chemother. Pharmacol.* **2012**, *69*, 679–689. [[CrossRef](#)] [[PubMed](#)]
57. Zupancic, S.; Kocbek, P.; Kristl, J.; Zariwala, M.; Renshaw, D.; Gul, M.O.; Elsaid, Z.; Taylor, K.; Somavarapu, S. Formulation and characterization of curcumin loaded DQAsomes for pulmonary delivery. In Proceedings of the 10th International Conference and Workshop on Biological Barriers, Saarbrücken, Germany, 16–21 February 2014.
58. Briskey, D.; Sax, A.; Mallard, A.R.; Rao, A. Increased bioavailability of curcumin using a novel dispersion technology system (LipiSpense®). *Eur. J. Nutr.* **2019**, *58*, 2087–2097. [[CrossRef](#)]
59. Faul, F.; Erdfelder, E.; Lang, A.-G.; Buchner, A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav. Res. Methods* **2007**, *39*, 175–191. [[CrossRef](#)]
60. Faul, F.; Erdfelder, E.; Buchner, A.; Lang, A.-G. Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Behav. Res. Methods* **2009**, *41*, 1149–1160. [[CrossRef](#)]
61. Study Randomizer. Available online: <https://app.studyrandomizer.com/> (accessed on 1 October 2019).
62. Dooley, J.; Worwood, M. *Guidelines on Diagnosis and Therapy: Genetic Haemochromatosis*; British Committee for Standards in Haematology: London, UK, 2000; pp. 1–33.
63. Fitzsimons, E.J.; Cullis, J.O.; Thomas, D.; Tsochatzis, E.; Griffiths, W.J. Diagnosis and therapy of genetic haemochromatosis (review and 2017 update). *Br. J. Haematol.* **2018**, *181*, 293–303. [[CrossRef](#)]
64. World Health Organization. *Nutritional Anaemias: Tools for Effective Prevention and Control*; World Health Organization: Geneva, Switzerland, 2017.
65. Punnonen, K.; Irjala, K.; Rajamäki, A. Serum Transferrin Receptor and Its Ratio to Serum Ferritin in the Diagnosis of Iron Deficiency. *Blood* **1997**, *89*, 1052–1057. [[CrossRef](#)]
66. Verdon, F.; Burnand, B.; Stubi, C.-L.F.; Bonard, C.; Graff, M.; Michaud, A.; Bischoff, T.; De Vevey, M.; Studer, J.-P.; Herzig, L.; et al. Iron supplementation for unexplained fatigue in non-anaemic women: Double blind randomised placebo controlled trial. *BMJ* **2003**, *326*, 1124. [[CrossRef](#)]
67. Koulaouzidis, A.; Cottier, R.; Bhat, S.; Said, E.; Linaker, B.D.; Saeed, A.A. A ferritin level >50 µg/L is frequently consistent with iron deficiency. *Eur. J. Intern. Med.* **2009**, *20*, 168–170. [[CrossRef](#)]
68. Vaucher, P.; Druais, P.-L.; Waldvogel, S.; Favrat, B. Effect of iron supplementation on fatigue in nonanemic menstruating women with low ferritin: A randomized controlled trial. *Can. Med. Assoc. J.* **2012**, *184*, 1247–1254. [[CrossRef](#)] [[PubMed](#)]
69. Peyrin-Biroulet, L.; Williet, N.; Cacoub, P. Guidelines on the diagnosis and treatment of iron deficiency across indications: A systematic review. *Am. J. Clin. Nutr.* **2015**, *102*, 1585–1594. [[CrossRef](#)] [[PubMed](#)]
70. Blood Pressure Association UK. Blood Pressure Chart [Bloodpressureuk.org](http://www.bloodpressureuk.org). 2008. Available online: <http://www.bloodpressureuk.org/BloodPressureandyou/Thebasics/Bloodpressurechart> (accessed on 5 May 2020).
71. Williams, B.; Mancia, G.; Spiering, W.; Rosei, E.A.; Azizi, M.; Burnier, M.; Clement, D.L.; Coca, A.; De Simone, G.; Dominiczak, A.; et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension. *J. Hypertens.* **2018**, *36*, 1953–2041. [[CrossRef](#)] [[PubMed](#)]
72. Goddard, A.F.; James, M.W.; McIntyre, A.S.; Scott, B.B. Guidelines for the management of iron deficiency anaemia. *Gut* **2011**, *60*, 1309–1316. [[CrossRef](#)] [[PubMed](#)]
73. British National Formulary. Ferrous Sulfate [bnf.nice.org.uk](https://bnf.nice.org.uk/drug/ferrous-sulfate.html). 2020. Available online: <https://bnf.nice.org.uk/drug/ferrous-sulfate.html> (accessed on 8 May 2020).
74. Food and Drug Administration (FDA) Center for Food Safety and Applied Nutrition. *Guidance for Industry: Food Labeling Guide*; Food and Drug Administration: College Park, MD, USA, 2013.
75. Coudène, P.; Marson, B.; Badiou, S.; Flavier, S.; Anelli, S.; Cristol, J.; Dupuy, A.M. Evaluation of the ABX Pentra 400: A newly available clinical chemistry analyser. *Clin. Chem. Lab. Med.* **2005**, *43*, 782–792. [[CrossRef](#)]
76. Simó, J.M.; Joven, J.; Clivillé, X.; Sans, T. Automated latex agglutination immunoassay of serum ferritin with a centrifugal analyzer. *Clin. Chem.* **1994**, *40*, 625–629. [[CrossRef](#)]

77. Tolkien, Z.; Stecher, L.; Mander, A.P.; Pereira, D.; Powell, J.J. Ferrous Sulfate Supplementation Causes Significant Gastrointestinal Side-Effects in Adults: A Systematic Review and Meta-Analysis. *PLoS ONE* **2015**, *10*, e0117383. [[CrossRef](#)]
78. World Health Organization (WHO). *Serum Ferritin Concentrations for the Assessment of Iron Status and Iron Deficiency in Populations*; World Health Organization: Geneva, Switzerland, 2011.
79. Mast, A.E.; Blinder, M.A.; Gronowski, A.M.; Chumley, C.; Scott, M.G. Clinical utility of the soluble transferrin receptor and comparison with serum ferritin in several populations. *Clin. Chem.* **1998**, *44*, 45–51. [[CrossRef](#)]
80. Clénin, G.E. The treatment of iron deficiency without anaemia (in otherwise healthy persons). *Swiss Med. Wkly.* **2017**, *147*, w14434.
81. Soppi, E. Iron Deficiency without Anemia—Common, Important, Neglected. *Clin. Case Rep. Rev.* **2019**, *5*, 1–7.
82. Cook, J.D.; Lipschitz, D.A.; Miles, L.E.M.; Finch, C.A. Serum ferritin as a measure of iron stores in normal subjects. *Am. J. Clin. Nutr.* **1974**, *27*, 681–687. [[CrossRef](#)] [[PubMed](#)]
83. Institute of Medicine (US) Panel on Micronutrients. *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*; National Academies Press: Washington, DC, USA, 2001.
84. Manoguerra, A.S.; Erdman, A.R.; Booze, L.L.; Christianson, G.; Wax, P.M.; Scharman, E.J.; Woolf, A.D.; Chyka, P.A.; Keyes, D.C.; Olson, K.R.; et al. Iron Ingestion: An Evidence-Based Consensus Guideline for Out-of-Hospital Management. *Clin. Toxicol.* **2005**, *43*, 553–570. [[CrossRef](#)] [[PubMed](#)]
85. Barbara, R.; Belletti, D.; Pederzoli, F.; Masoni, M.; Keller, J.; Ballestrazzi, A.; Vandelli, M.; Tosi, G.; Grabrucker, A. Novel Curcumin loaded nanoparticles engineered for Blood-Brain Barrier crossing and able to disrupt Abeta aggregates. *Int. J. Pharm.* **2017**, *526*, 413–424. [[CrossRef](#)] [[PubMed](#)]
86. Hoppe, J.B.; Coradini, K.; Frozza, R.L.; Oliveira, C.M.; Meneghetti, A.B.; Bernardi, A.; Pires, E.S.; Beck, R.C.; Salbego, C.G. Free and nanoencapsulated curcumin suppress β -amyloid-induced cognitive impairments in rats: Involvement of BDNF and Akt/GSK-3 β signaling pathway. *Neurobiol. Learn. Mem.* **2013**, *106*, 134–144. [[CrossRef](#)]
87. Navaratna, D.; Guo, S.-Z.; Hayakawa, K.; Wang, X.; Gerhardinger, C.; Lo, E.H. Decreased Cerebrovascular Brain-Derived Neurotrophic Factor—Mediated Neuroprotection in the Diabetic Brain. *Diabetes* **2011**, *60*, 1789–1796. [[CrossRef](#)]
88. Moretti, D.; Goede, J.S.; Zeder, C.; Jiskra, M.; Chatzinakou, V.; Tjalsma, H.; Melse-Boonstra, A.; Brittenham, G.; Swinkels, R.W.; Zimmermann, M.B. Oral iron supplements increase hepcidin and decrease iron absorption from daily or twice-daily doses in iron-depleted young women. *Blood* **2015**, *126*, 1981–1989. [[CrossRef](#)]
89. Nemeth, E.; Ganz, T. Regulation of Iron Metabolism by Hepcidin. *Annu. Rev. Nutr.* **2006**, *26*, 323–342. [[CrossRef](#)]
90. Rossi, E. Hepcidin—The iron regulatory hormone. *Clin. Biochem. Rev.* **2005**, *26*, 47.
91. Sangkhae, V.; Nemeth, E. Regulation of the Iron Homeostatic Hormone Hepcidin. *Adv. Nutr.* **2017**, *8*, 126–136. [[CrossRef](#)]
92. Stoffel, N.U.; Cercamondi, C.I.; Brittenham, G.; Zeder, C.; Geurts-Moespot, A.J.; Swinkels, D.W.; Moretti, D.; Zimmermann, M. Iron absorption from oral iron supplements given on consecutive versus alternate days and as single morning doses versus twice-daily split dosing in iron-depleted women: Two open-label, randomised controlled trials. *Lancet Haematol.* **2017**, *4*, e524–e533. [[CrossRef](#)]
93. Granick, S. Protein Apoferritin and Ferritin in Iron Feeding and Absorption. *Science* **1946**, *103*, 107. [[CrossRef](#)] [[PubMed](#)]
94. Fuqua, B.K.; Vulpe, C.D.; Anderson, G.J. Intestinal iron absorption. *J. Trace Elem. Med. Biol.* **2012**, *26*, 115–119. [[CrossRef](#)] [[PubMed](#)]
95. Peña-Rosas, J.P.; De-Regil, L.M.; Malave, H.G.; Flores-Urrutia, M.C.; Dowswell, T. Intermittent oral iron supplementation during pregnancy. *Cochrane Database Syst. Rev.* **2015**, *2015*, CD009997. [[CrossRef](#)] [[PubMed](#)]
96. Galy, B.; Ferring-Appel, D.; Becker, C.; Gretz, N.; Gröne, H.-J.; Schümann, K.; Hentze, M.W. Iron Regulatory Proteins Control a Mucosal Block to Intestinal Iron Absorption. *Cell Rep.* **2013**, *3*, 844–857. [[CrossRef](#)]
97. Cegarra, L.; Colins, A.; Gerdtzen, Z.P.; Nuñez, M.T.; Salgado, J.C. Mathematical modeling of the relocation of the divalent metal transporter DMT1 in the intestinal iron absorption process. *PLoS ONE* **2019**, *14*, e0218123. [[CrossRef](#)]

98. Stoffel, N.U.; Zeder, C.; Brittenham, G.M.; Moretti, D.; Zimmermann, M.B. Iron absorption from supplements is greater with alternate day than with consecutive day dosing in iron-deficient anemic women. *Haematologica* **2019**, *105*, 1232–1239. [[CrossRef](#)]
99. Lund, E.K.; Wharf, S.G.; Fairweather-Tait, S.J.; Johnson, I. Oral ferrous sulfate supplements increase the free radical—Generating capacity of feces from healthy volunteers. *Am. J. Clin. Nutr.* **1999**, *69*, 250–255. [[CrossRef](#)]
100. Lund, E.K.; Fairweather-Tait, S.J.; Wharf, S.G.; Johnson, I. Chronic exposure to high levels of dietary iron fortification increases lipid peroxidation in the mucosa of the rat large intestine. *J. Nutr.* **2001**, *131*, 2928–2931. [[CrossRef](#)]
101. Jaeggi, T.; Kortman, G.A.M.; Moretti, D.; Chassard, C.; Holding, P.; Dostal, A.; Boekhorst, J.; Timmerman, H.M.; Swinkels, R.W.; Tjalsma, H.; et al. Iron fortification adversely affects the gut microbiome, increases pathogen abundance and induces intestinal inflammation in Kenyan infants. *Gut* **2015**, *64*, 731–742. [[CrossRef](#)]
102. Paganini, D.; Uyoga, M.A.; Kortman, G.A.M.; Cercamondi, C.I.; Moretti, D.; Barth-Jaeggi, T.; Schwab, C.; Boekhorst, J.; Timmerman, H.M.; Lacroix, C.; et al. Prebiotic galacto-oligosaccharides mitigate the adverse effects of iron fortification on the gut microbiome: A randomised controlled study in Kenyan infants. *Gut* **2017**, *66*, 1956–1967. [[CrossRef](#)]
103. Di Meo, F.; Margarucci, S.; Galderisi, U.; Crispi, S.; Peluso, G. Curcumin, Gut Microbiota, and Neuroprotection. *Nutrients* **2019**, *11*, 2426. [[CrossRef](#)] [[PubMed](#)]
104. Hibberd, M.C.; Wu, M.; Rodionov, D.A.; Li, X.; Cheng, J.; Griffin, N.W.; Barratt, M.J.; Giannone, R.J.; Hettich, R.L.; Osterman, A.L.; et al. The effects of micronutrient deficiencies on bacterial species from the human gut microbiota. *Sci. Transl. Med.* **2017**, *9*, eaal4069. [[CrossRef](#)] [[PubMed](#)]
105. Pivac, N.; Kim, B.; Nedic, G.; Joo, Y.H.; Kozaric-Kovacic, D.; Hong, J.P.; Mück-Šeler, D. Ethnic Differences in Brain-derived Neurotrophic Factor Val66Met Polymorphism in Croatian and Korean Healthy Participants. *Croat. Med. J.* **2009**, *50*, 43–48. [[CrossRef](#)] [[PubMed](#)]
106. Bus, B.A.; Molendijk, M.; Penninx, B.; Buitelaar, J.; Kenis, G.; Prickaerts, J.; Elzinga, B.; Voshaar, R.O. Determinants of serum brain-derived neurotrophic factor. *Psychoneuroendocrinology* **2011**, *36*, 228–239. [[CrossRef](#)]
107. Naegelin, Y.; Dingsdale, H.; Säuberli, K.; Schädelin, S.; Kappos, L.; Barde, Y.-A. Measuring and Validating the Levels of Brain-Derived Neurotrophic Factor in Human Serum. *Eneuro* **2018**, *5*, 1–9. [[CrossRef](#)]
108. Polacchini, A.; Metelli, G.; Francavilla, R.; Baj, G.; Florean, M.; Mascaretti, L.G.; Tongiorgi, E. A method for reproducible measurements of serum BDNF: Comparison of the performance of six commercial assays. *Sci. Rep.* **2015**, *5*, 17989. [[CrossRef](#)]
109. Matrisciano, F.; Bonaccorso, S.; Ricciardi, A.; Scaccianoce, S.; Panaccione, I.; Wang, L.; Ruberto, A.; Tatarelli, R.; Nicoletti, F.; Girardi, P.; et al. Changes in BDNF serum levels in patients with major depression disorder (MDD) after 6 months treatment with sertraline, escitalopram, or venlafaxine. *J. Psychiatr. Res.* **2009**, *43*, 247–254. [[CrossRef](#)]
110. Mu, J.-S.; Li, W.-P.; Yao, Z.-B.; Zhou, X.-F. Deprivation of endogenous brain-derived neurotrophic factor results in impairment of spatial learning and memory in adult rats. *Brain Res.* **1999**, *835*, 259–265. [[CrossRef](#)]
111. Tapia-Arancibia, L.; Aliaga, E.; Silhol, M.; Arancibia, S. New insights into brain BDNF function in normal aging and Alzheimer disease. *Brain Res. Rev.* **2008**, *59*, 201–220. [[CrossRef](#)]
112. Piepmeier, A.; Etnier, J.L. Brain-derived neurotrophic factor (BDNF) as a potential mechanism of the effects of acute exercise on cognitive performance. *J. Sport Health Sci.* **2015**, *4*, 14–23. [[CrossRef](#)]
113. Miranda, M.; Morici, J.F.; Zanoni, M.B.; Bekinschtein, P. Brain-Derived Neurotrophic Factor: A Key Molecule for Memory in the Healthy and the Pathological Brain. *Front. Cell. Neurosci.* **2019**, *13*, 363. [[CrossRef](#)] [[PubMed](#)]
114. Shimada, H.; Makizako, H.; Doi, T.; Yoshida, D.; Tsutsumimoto, K.; Anan, Y.; Uemura, K.; Lee, S.; Park, H.; Suzuki, T. A Large, Cross-Sectional Observational Study of Serum BDNF, Cognitive Function, and Mild Cognitive Impairment in the Elderly. *Front. Aging Neurosci.* **2014**, *6*, 69. [[CrossRef](#)] [[PubMed](#)]

